
In vitro* evaluation of *Croton bonplandianum* Baill. as potential antitumor properties using *Agrobacterium tumefaciens

Islam, M.S., Rahman, M.M., Rahman, M.A., Qayum, M.A. and Alam, M.F.*

Biotechnology and Microbiology Laboratory, Department of Botany, University of Rajshahi, Rajshahi-6205, Bangladesh.

Islam, M.S., Rahman, M.M., Rahman, M.A., Qayum, M.A. and Alam, M.F. (2010). *In vitro* evaluation of *Croton bonplandianum* Baill. as potential antitumor properties using *Agrobacterium tumefaciens*. Journal of Agricultural Technology 6(1): 79-86.

Antitumor properties of twigs extract of *Croton bonplandianum* Baill. were proven using potato disc and radish seed bioassays. Tumor formation ability of *Agrobacterium* was distinctly inhibited on potato disc in presence of methanol extract. Significant variation on the ability of tumor formation was observed among the tested strains of *Agrobacterium* when treated with different concentrations (10, 100 and 1,000 ppm) of plant extracts. Tumor inhibition was increased with the increasing of concentrations of plant extract. Decreasing the root length and percentage of seed germination during radish seed bioassays further confirmed the antitumor properties of *Croton bonplandianum*. In conclusion it may be recommended that the bioactive compound of this plant may play an important role in developing antitumor drugs for human beings, as there is a similarity between human and plant tumor formation mechanism.

Key words: Antitumor properties, *Croton bonplandianum*, potato disc bioassay, *Agrobacterium*

Introduction

Cancer is now serious health problems in human beings both in developed and developing countries. Some conventional systems such as surgery, chemotherapy, radiation therapy, immunotherapy, monoclonal antibody therapy or other methods are being used for Cancer treatment. Most of the agents have been revealed as mutagenic and/or carcinogenic, and are highly toxic, not only for cancer but also for normal cells (Fatber, 1968). Due to the toxic and adverse side effects of synthetic medicines being observed round the globe, herbal medicine has made a comeback to improve the fulfillment of our present and future health needs (Harun-ur-Rashid *et al.*, 2002). Several plant-derived compounds have been approved as anti-cancer drugs i.e. vinblastine,

*Corresponding author: M. Firoz Alam; e-mail: falambt@yahoo.com

vincristine, etoposide, teniposide, taxol, taxotere, topotecan and irinotecan, just to name a few (Syrovets and Laumonnier, 2009).

Crown gall is a neoplastic disease of plants caused by *Agrobacterium tumefaciens* (Kahl and Schell, 1982; Lippincott and Lippincott, 1975) which occurs in more than 60 families of dicotyledons and many gymnosperms (Galsky and Wilsey, 1980). The Ti-plasmid causes the plant's cells to multiply rapidly without going through apoptosis, resulting in tumor formation similar in nucleic acid content and histology to human and animal cancers (McLaughlin, 1991; Agrios, 1997). The potato disc assay demonstrates the inhibition of tumor formation on potato discs; materials (e.g. plant extracts) that inhibit these plant tumors have a high predictability of showing activity against the P388 (3PS) leukemia in mice (Ferrigni *et al.*, 1982). Development of a simple antitumor prescreen using a convenient and inexpensive plant tumor assay systems can offer numerous advantages as alternatives to extensive animal testing in the search for new anticancer drugs.

Croton bonplandianum Baill. is native to the southern Bolivia, Paraguay, Southwestern Brazil, and Northern Argentina of South America (Chakrabarty and Balakrishnan, 1992). *Croton* is rich in secondary metabolites including alkaloids and terpenoids (Rizk, 1987), the latter including irritant co-carcinogenic phorbol esters (Phillipson, 1995). Diterpene resins found in many species of *Croton* have been used experimentally in studies of tumor initiation and conceivably prove to be useful in cancer therapy (Sharma, 2009). Considering its tremendous important in antitumor potential, the present study was undertaken to evaluate their antitumor and phytotoxicity by utilization of potato disc and radish seed bioassay.

Materials and methods

Plant material

Croton bonplandianum Baill. was collected from different places of Rajshahi University Campus. Taxonomic identity of this plant was confirmed by Mr. A.H.M. Mahbubur Rahman, Assistant Professor, Department of Botany, Rajshahi University, Rajshahi-6205, Bangladesh. Twigs were used for extraction.

Preparation of extracts

Extraction procedure was carried out according to Ahmad and Beg (2001) with some modification. Twigs were rinsed well with tap and distilled water (DW) and kept under shade still drying. Dried material coarsely powdered using mortar and pestle followed by oven dry and further reduced to powder

using an electric blender (Nokia, Osaka-Japan) and stored in air tight glass container. Powder (50 g) was then dissolved in methanol by allowing to sediment at room temperature (27-30 °C) for 7 days with occasional shaking. For filtrating the material, Teton cloth and Whatman No.1 filter paper were used, respectively. The filtrates were then transferred into glass beaker and dried into semisolid material using water bath (4 holes analogue, Thermostatic water bath, China). Particular concentration (10 ppm, 100 ppm and 1,000 ppm) of the plant extract was prepared using methanol for antitumor activity test, and 250 mg/ml was prepared for antibacterial assay.

Antitumor potato disc assay

Antitumor test of plant extracts was performed according to standard potato disc bioassay (Hussain *et al.*, 2007). Three *A. tumefaciens* strains named AtTa0112, AtAc0114 and AtSl0105 (isolated and identified in our laboratory) were cultured on Luria-Bertani (LB) agar and then transferred into LB broth and incubated 48 hours. Six to seven loops of broth cultures were transferred into test tube containing 10 ml phosphate buffered saline (PBS; pH 7.2). The following proportion was used for antitumor activity test: 600 µl test extract + 150 µl Double Distilled Water (DDW) + 750 µl *A. tumefaciens* in PBS. Camptothecin was used as positive control replacing test extracts.

Red skinned potatoes (*Solanum tuberosum* L.) were collected from local market and thoroughly washed with tap and DW. For surface sterilization, 0.1% HgCl₂ solution was used. Potato tubers were cut into 8 mm diameter in size cylindrical pieces using cork borer and transferred into DDW containing conical flask. After washing, the cylindrical segments were cut into 5 mm × 8 mm disc and placed onto agar (15 g/l) plates (10 discs per plate). A 50 µl of appropriate inoculum were placed on the surface of each potato disc. The plates were sealed with parafilm and incubated at room temperature at 27 - 30 °C for 21 days. After 21 days discs were stained with Lugol's solutions (10% KI, 5% I₂) and tumors were counted under a stereo microscope. The experiment was carried out in sterilized condition and repeated three times. Percent inhibition of tumors was calculated (McLaughlin, 1991; McLaughlin *et al.*, 1993; McLaughlin and Rogers, 1998). More than 20% tumor inhibition is considered significant (Ferrigni *et al.*, 1982). Data were analyzed using ANOVA.

Antibacterial assay

The disc diffusion assay (Kirby-Bauer Method) was used to screen for antibacterial activity of tested plant extract (Bauer *et al.*, 1966; Barry, 1980). Methanol was used as negative control and erythromycin (30 µg/ml),

carbenicillin (100 µg/ml) and chloramphenicol (30 µg/ml) were used as positive control. Discs (Whatman No.1 filter paper) were impregnated with 10 µl of the extract and antibiotics followed by air dried, and then placed on seeded LB agar plates. 20 µl standard bacterial cultures (48 hours incubated) were used for spreading LB agar plates. Plates were then incubated at 28-30 °C for 24 hours. Antibacterial activity was evaluated by measurement of diameter zones of inhibition (mm) against *A. tumefaciens* strains. Each assay was carried out in triplicates.

Radish seed phytotoxicity assay

Radish seed phytotoxicity assay was conducted according to Turker and Camper (2002). Two different determinations: 1) root length and 2) percent seed germination were performed. For root length determination, two different concentrations (1,000 ppm and 10,000 ppm) of the extract were used. Five ml of each concentration of the extract was applied on filter paper in Petri plates. After evaporation of methanol, 5 ml DW and sterilized (using 0.1% HgCl₂) 20 radish seeds were placed in each Petri plate. Petri plates were sealed with parafilm and incubated at 23 ± 2 °C. DDW was used as a negative control. Root length was measured after 1, 3 and 5 days of interval. This experiment was repeated three times.

For seed germination, two different concentrations (1,000 and 7,500 ppm) of the extract were used. This part of the experiment is similar to that of earlier experiment except for the extract concentrations and number of seeds. Here, 100 radish seeds were placed on each Petri plate. Germinated seeds were counted every day up to 5 days after treatment. Experiment was repeated three times.

Results

Antitumor potato disc assay

Three strains of tumor producing *A. tumefaciens* showed significantly different response from each others in all tested concentrations of the plant extracts. Similarly, the extract concentration also significantly differed in tumor inhibition. Maximum and minimum tumor inhibition were observed against the strain AtSI0105 (16.92-40.76%) and AtTa0112 (9.83-37.70%), respectively. No significant tumor inhibition was observed at 10 ppm against the studied strains. Strain AtSI0105 was more prominent for producing tumor (8.6 ± 0.23) compared to strains AtAc0114 (7.2 ± 0.20) and AtTa0112 (6.1 ± 0.25) (Fig.1, 2).




Fig.1. Comparative graphical presentation of different concentrations of methanol extract of *C. bonplandianum* and studied three *A. tumefaciens* strains on crown gall tumors formation in potato discs.




Fig. 2. Effect of methanol extract of *C. bonplandianum* on crown gall tumors formation by *A. tumefaciens* strains on potato discs. Data were recorded after 21 days of treatment. A1, B1 and C1 showing normal tumor development in absence of plant extracts (control). A2, B2 and C2 showing reduction of number of tumor development in presence of 10 ppm plant extract. A3, B3 and C3 showing reduction of tumor formation in presence of 1,000 ppm plant extract.

Antibacterial assay

The results of antitumor potential of the plant extract but not on *Agrobacterium* viability, antibacterial assay were justified before antitumor activity test. A high concentration (250 mg/ml) of the plant extract was used for antibacterial assay and no inhibition zone was recorded. Similar results were observed in antibiotic treatment.

Radish seed phytotoxicity assay

Root length and seeds germination were actively inhibited by the extracts at both concentration (1,000 ppm and 10,000 ppm) (Fig. 3, 4). On the fifth day, root length 14.56 ± 0.34 and 7.16 ± 0.37 were recorded at 1,000 ppm and 10,000 ppm, respectively. On the other hand, seed germinations (45.66 ± 0.120 and 26.66 ± 0.88) were recorded at 1,000 ppm and 7,500 ppm, respectively.




Fig. 3. Histogram showing radish seed phytotoxicity assay on root length at two concentrations of methanol extract of *C. bonplandianum* including control.




Fig. 4. Graph showing radish seed phytotoxicity assay on percentage of seed germination at two concentrations of methanol extract of *C. bonplandianum* including control.

Discussions

Antitumor potato disc assay is a valuable tool that indicates antitumor activity of test compound by their inhibition crown galls formation that was induced in wounded potato tissues by *A. tumefaciens*. This bioassay was a sensitive, bench-top antitumor assay for chemicals that disrupt the cell cycle (mitosis, S phase, etc.) regardless of their mode of action (Coker *et al.*, 2003). Several scientists have been used these methods over the past 15 years, and they appear to be adaptable to the purpose of standardization or quality control of bioactive compounds in such heterogeneous botanicals (Jerry and Lingling, 1998).

C. bonplandianum plays an important in antitumor potential, the methanol extract was evaluated for antitumor activity with phytotoxic analysis. It was shown that tumor formation was observed when *Agrobacterium* strains alive on living potato disc. The potato discs were often damaged due to the contamination and other physiological factors when there was no tumor formation. Thus successful attachment of *Agrobacterium* on living potato disc is needed for antitumor test of plant extracts.

The results of antitumor activity and antibacterial showed that a high concentration 250 mg/ml of the plant extract had no inhibition zone. This indicated that there was no effect of plant extract on the viability of *A. tumefaciens* also as reported by Hussain *et al.* (2007). Erythromycin, carbenicillin and chloramphenicol were tested for further confirmation and similar results were found. These results revealed that tumor formation was decreased only for the bioactive compound present in plant extract.

This study root length and seed germination were actively inhibited by the plant extract at 1,000 ppm., as a result supported by Inayatullah *et al.* (2007). The maximum inhibition was observed at 10,000 ppm as reported earlier by Turker and Camper (2002).

References

- Agrios, G.N. (1997). Plant diseases caused by prokaryotes: bacteria and mollicutes. Plant Pathology. Academic Press, San Diego, USA.
- Ahmad, I. and Beg, Z. (2001). Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. Journal of Ethnopharmacology 74: 87-91.
- Barry, A.L. (1980). Procedure for testing antimicrobial agent in agar media. In: Lorian V (ed) Antibiotica in laboratory medicines, Willims & Wilkins Co. Baltimore, USA.
- Bauer, A.W., Kibry, W.M.M., Sherris, J.C. and Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disc method. American Journal of Clinical Pathology 45: 493-496.

- Chakrabarty, T. and Balakrishnan, N.P. (1992). A revision of *Croton* L. (Euphorbiaceae) for Indian subcontinent. *Bulletin of the Botanical Survey of India* 34(1-4): 1-88.
- Coker, P.S., Radecke, J., Guy, C. and Camper, N.D. (2003). Potato tumor induction assay: a multiple mode of action drug assay. *Phytomedicine* 10: 133-138.
- Fatber, B. (1968). Biochemistry of Carcinogenesis. *Cancer Research* 28: 2338-2349.
- Ferrigni, N.R., Putnam, J.E., Anderson, B., Jacobsen, L.B., Nichols, D.E., Moore, D.S. and McLaughlin, J.L. (1982). Modification and evaluation of the potato disc assay and antitumor screening of Euphorbiaceae seeds. *Journal of Natural Products* 45: 679-686.
- Galsky, A.B. and Wilsey, J.P. (1980). Crown-gall tumor disc bioassay: a possible aid in the detection of compounds with antitumor activity. *Plant Physiology* 65: 184-185.
- Harun-ur-Rashid, M., Gafur, M.A., Sadik, G.M. and Rahman, M.A.A. (2002). Biological activities of a new acrylamide derivative from *Ipomoea turpithum*. *Pakistan Journal of Biological Science* 5(9): 968-969.
- Hussain, A., Zia, M. and Mirza, B. (2007). Cytotoxic and antitumor potential of *Fagonia cretica* L. *Turkish Journal of Biology* 31: 19-24.
- Jerry, L.M. and Lingling, L.R. (1998). The use of biological assays to evaluate botanicals. *Drug Information Journal* 32: 513-524.
- Kahl, G. and Schell, J.S. (1982). *Molecular Biology of Plant Tumors*. Academic Press, New York.
- Lippincott, J.A. and Lippincott, B.B. (1975). The genus *Agrobacterium* and plant tumorigenesis. *Annual Review of Microbiology* 29: 377-405.
- McLaughlin, J.L. (1991). Crown gall tumors on potato discs and bine shrimp lethality: Two single bioassays for plant screening and fractionation. In: Hostettmann K (ed) *Methods in Plant Biochemistry*, Vol 6. Academic Press, London. p. 1-31.
- McLaughlin, J.L. and Rogers, L.L. (1998). The use of biological assays to evaluate botanicals. *Drug Information Journal* 32: 513-524.
- McLaughlin, J.L., Chang, C.J. and Smith, D.L. (1993). Simple bench-top bioassay (brine shrimp and potato discs) for the discovery of plant antitumor compounds. In: Kinghorn, A.D., Balandrin, M.F. (eds) *Human Medicinal Agents from Plants*. ACS Symposium. Series 534. Washington, D.C.: American Chemical Society. P. 112-137.
- Phillipson, J.D. (1995). A matter of some sensitivity. *Phytochemistry* 38: 1319-1343.
- Rizk, A.F.M. (1987). The chemical constituents and economic plants of the Euphorbiaceae. *Botanical Journal of the Linnean Society* 94: 293-326.
- Sharma, H.M. (2009). [Online] Available: <http://zolengthe.blog.co.in/2008/06/27/the-ethnomedicinal-use-of-croton/>.
- Syrovets, T. and Laumonier, Y. (2009). [Online]. Available: http://www.uni-ulm.de/klinik/nhk/dept/research/projekte_Syrovets_neu.htm.

(Received 20 April 2009; accepted 25 October 2009)