
Genetic Diversity of *Elaeagnus Latifolia* L. as Revealed by Random Amplified Polymorphic DNA (RAPD) Markers

Yingthongchai, P.¹, Naphrom, D.² and Smitamana, P.^{1*}

¹Plant Biotechnology Program, Graduate School, Chiang Mai University, Chiang Mai, 50200, Thailand, ²Department of Plant Science and Natural Resources, Faculty of Agriculture, Chiang Mai University, Chiang Mai, 50200, Thailand.

Yingthongchai, P., Naphrom, D. and Smitamana, P. (2014). Genetic diversity of *Elaeagnus latifolia* L. as revealed by random amplified polymorphic DNA (RAPD) markers. International Journal of Agricultural Technology 10(1):177-187.

Abstract *Elaeagnus latifolia* L., an endemic fruit plant, is found in the upper north of Thailand. Sixty accessions were collected from seven locations: Chiang Mai, Chiang Rai, Lamphun, Lampang, Nan, Phayao and Mae Hong Son. Randomly amplified polymorphic DNA (RAPD) markers were used to identify genetic diversity of *E. latifolia*. Seven primers were screened from 50 primers and provided a total of 178 DNA fragments of which 168 (94.38 %) were polymorphic. The genetic similarity matrix was calculated using the Jaccard similarity coefficient, and SAHN procedure was used for Dendrogram clustering. Jaccard's similarity coefficient (GSj) ranged from 0.38-0.71 and the cophenetic correlation of the similarity matrix (r) was 0.73. Fifty eight out of 60 accessions could be distinguished where as only two accessions; N55 and N72 appeared to be close to each other, with the genetic similarity coefficient of 0.71. The results indicated that the RAPD method is sufficiently informative and powerful to assess the genetic diversity in *E. latifolia*.

Keywords: *Elaeagnus latifolia* L., DNA fingerprinting, Genetic diversity, RAPD markers

Introduction

The plants in the genus *Elaeagnus* (Family Elaeagnaceae) are distributed widely from the northern regions of Asia to the Himalayas, and different parts of Europe and North America. In India they grow up to an altitude of 2300 m in the Himalayas (Ahmadiani *et al.*, 2000; Bhattacharyya and Johri, 1998). Different species of *Elaeagnus* are cultivated as deciduous or evergreen ornamental shrubs. This genus is known to contain mono- and disaccharides in the leaves, fructose and L-ascorbic acid in the fruits, fatty acids and phytosterols in the seed, leaves and stems and carotenoids in the fruits (Sakamura and Suga, 1987).

One species of this genus, *Elaeagnus latifolia* L., is an endemic fruit plant mostly found in the upper north of Thailand. The plant has long been known by

* **Corresponding author:** Smitamana, P.; **Email:** psporn@gmail.com

the local people and available in local markets where fruit were sold for about 10-15 Baht/kg. It is a climbing shrub rarely a tree; a scandent shrub with rusty shiny scales that are often thorny or spiny. Fruit is considered to be a very rich source of vitamins and minerals and other bioactive compounds. It is also a good source of essential fatty acids, which is unusual in other fruit trees. The tree is a great potential crop because of its good adaptability and toleration to different habitats, early bearing, high yield and various uses of ripe fruit. Fruit are eaten raw and used for making chutney, fruit pulp was used for making jam, jelly and refreshing drinks. The fruit of *E. latifolia* is being evaluated as a food that is capable of reducing the incidence of cancer and also as a means of halting or reversing the growth of cancers (Patel *et al.*, 2008; Sundriyal and Sundriyal, 2003; Sundriyal and Sundriyal, 2005). *E. latifolia* is an actinorhizal plant (forms a symbiosis with the nitrogen-fixing bacterium *Frankia*) that has been studied in Kumuan Himalaya for its ability to restore degraded lands, and its leaves provide good source of fodder (Bargali, 2011).

In recent years, populations of *E. latifolia* have rapidly declined and the plant has disappeared from many locations of Thailand. The risk of the occurrence can be caused by the plant as endemic species and the fruits do not well known in the markets, therefore it is left out and only seasonal fruit picking without further proper propagation has been done. Moreover, the areas are replanted with the other more economically important crops. Therefore, characterization of the genetic diversity of *E. latifolia* is important for sustainable genetic conservation and increased use of plant genetic resources

Materials and methods

Plant materials

Sixty accessions of *E. latifolia*, were collected from seven provinces in the upper north of Thailand, i.e. Chiang Mai (CM), Chiang Rai (CR), Mae Hong Son (MH), Nan (N), Phayao (PY), Lamphun (L) and Lampang (LP) for RAPD analysis (Fig. 1 and Table 1). Two to three young leaves were removed from each tree and stored in freezers at -20 °C. until use.

DNA extraction

Total genomic DNA was extracted from approximately 100 mg young leaves using DNeasy Plant Mini kit (Qiagen) following the protocol of the manufacturer. The quantity and quality of isolated total genomic DNA was determined using 0.8% agarose gel electrophoresis in 0.5X TBE buffer for

mobility relative to know concentration of EZ load Precision Molecular Mass Standard and diluted to uniform concentration (10 ng/μl) for RAPD analysis.

DNA amplification

Seven primers were screened from 50 primers) Operon Technology, USA). DNA amplification was carried out in 25 μl reactions consisting of pureTaq Ready-To-Go PCR Beads (GE Healthcare UK Limited, UK), 0.2 μM of each primer, 10 ng of template DNA and double-distilled water. The reaction was performed in Gradient Palm-Cycle™ (The Corbett Research Life Science, Australia). Amplification condition consisted of pre denaturation at 94 °C for 5 min, followed by 2 cycles of 94 °C for 60 sec, 38 °C for 7 sec, 72°C for 70 sec, and 55 cycles of 92 °C for 2 sec, 40 °C for 7 sec, 72 °C for 70 sec, followed by a final extension of 72 °C for 5 min. Amplification products were analyzed by electrophoresis at a constant voltage of 120 V for 2 h, in 1.5 % agarose gels with 0.5X TBE buffer. A 100 bp ladder plus (Fermentas, USA) was used to estimate the molecular size of the fragments. After electrophoresis the gel was stained with ethidium bromide (0.5 μg/ml) and the patterns were photographed using a gel documentation system.

Data analysis

The observed bands in the gel were evaluated based on the presence (1) or absence (0) of polymorphic fragments for each primer. Only data from intensely stained, unambiguous, clearly visible bands were used for statistical analysis. Cluster analysis was performed with NTSYS-pc. Version 2.2, a numerical taxonomy and multivariate analysis software package (Rohlf, 2005) using unweighted pair-group method with an arithmetic average (UPGMA). Genetic similarities between samples were measured by the Jaccard's similarity coefficient (GS_j) with SIMQUAL module. The distance coefficients were used to construct a dendrogram using UPGMA employing the Sequential Agglomerative Hierarchical and Nested (SAHN) algorithm. The goodness of fit of the clustering compared with the basic data matrix was tested by computing the cophenetic correlation coefficient using the normalized Mantel statistics Z test via the COPH and MXCOMP procedures. Polymorphic information content (PIC) was calculated for RAPD markers; $PIC = 1 - \sum P_{ij}^2$, where relative frequency of *j*th band for *i*th primer is summed across all the bands for the primer over all accessions (Li *et al.*, 2009).



Fig. 1. *Elaeagnus latifolia* L.

Table 1. Sixty accessions of *Elaeagnus latifolia* L. used in RAPD analysis

Accession No.	Accession code	Location	Accession No.	Accession code	Location
1.	L2	Lamphun	31.	H46	Chiang Mai
2.	L31	Lamphun	32.	MT12	Chiang Mai
3.	L42	Lamphun	33.	MT16	Chiang Mai
4.	L56	Lamphun	34.	D2	Chiang Mai
5.	L57	Lamphun	35.	CM80	Chiang Mai
6.	L59	Lamphun	36.	CM114	Chiang Mai
7.	L65	Lamphun	37.	N55	Nan
8.	L71	Lamphun	38.	N72	Nan
9.	L72	Lamphun	39.	N76	Nan
10.	LP 20	Lampang	40.	N99	Nan
11.	LP 53	Lampang	41.	N100	Nan
12.	LP 59	Lampang	42.	N101	Nan
13.	LP 61	Lampang	43.	N110	Nan
14.	LP 67	Lampang	44.	PY16	Phayao
15.	LP 68	Lampang	45.	PY52	Phayao
16.	LP 70	Lampang	46.	PY56	Phayao
17.	MH 5	Mae Hong Son	47.	PY62	Phayao
18.	MH 31	Mae Hong Son	48.	PY63	Phayao
19.	MH 49	Mae Hong Son	49.	PY68	Phayao
20.	MH 54	Mae Hong Son	50.	PY85	Phayao
21.	MH 60	Mae Hong Son	51.	CR29	Chiang Rai

22.	MH 63	Mae Hong Son	52.	CR66	Chiang Rai
23.	MH 70	Mae Hong Son	53.	CR70	Chiang Rai
24.	MH 72	Mae Hong Son	54.	CR90	Chiang Rai
25.	MH 74	Mae Hong Son	55.	CR98	Chiang Rai
26.	H 15	Chiang Mai	56.	CR103	Chiang Rai
27.	H19	Chiang Mai	57.	CR104	Chiang Rai
28.	H26	Chiang Mai	58.	CR108	Chiang Rai
29.	H31	Chiang Mai	59.	CR113	Chiang Rai
30.	H38	Chiang Mai	60.	CR119	Chiang Rai

Results

Out of 50 random decamer primers screened for polymorphism, 7 primers were selected for characterization of 60 accessions of *E. latifolia* (OPL-12, OPO-19, OPR-04, OPT-12, OPY-04, OPU-13, OPAJ-18 and OPC-19). Amplification of genomic DNA of 60 accessions using 7 primers for RAPD analysis gave the results shown in Table 2. The number of fragments produced by each primer ranged from 18 (OPT-12) to 31 (OPO-19 and OPR-04). A total of 178 DNA fragments were identified, 168 of them were polymorphic with an average of 24.00 polymorphic fragments per primer. The percentage of polymorphic bands (PPB) ranged from 86.36% (OPY-04) to a maximum of 100% (OPL-12 and OPO-19), with an average of 94.38 % polymorphism. Fig. 2 and Fig. 3 represent the extent of polymorphism observed among 60 accessions of *E. latifolia* as revealed by OPL-12 and OPO-19, respectively. The mean PIC value observed was 0.285 across all primers (Table 2).

The amplified RAPD bands were adequate to discriminate all accessions. The polymorphic bands were used to generate the similarity matrix. Jaccard's similarity coefficient or genetic similarity (GS_j) ranged from 0.38-0.71. Dendrogram based on UPGMA analysis grouped the 60 accessions into main clusters. The 60 accessions from RAPD analysis were grouped into one main cluster and a single accession, H38, formed a separate out group showing a low similarity coefficient (0.38) with other accessions (Fig. 4). Accessions within main cluster were grouped into five groups. The first group I comprised L2, L42 and PY68. The group II consisted of 16 accessions, 6 accessions collected from Lamphun (L), 6 accessions collected from Lampang (LP) and 4 accessions from Mae Hong Son (MH). Group III was comprised of 21 accessions, 7 accessions collected from Nan (N), 9 accessions collected from Chiang Rai (CR) and 5 accessions collected from Phayao (PY). Within Nan, N55 and N72 appeared to be closer to each other, with the genetic similarity

coefficient of 0.71. Their relatively high similarity may indicate that they developed from a common ancestor with similar fruit shape and taste. Group IV consisted of 13 accessions, 3 accessions collected from Mae Hong Son (MH) and 10 accessions collected from Chiang Mai (CM). The last group, group V comprised L59, MH61, MH63, LP59, CR29 and PY85. A cophenetic correlation tree of 60 accessions of *E. latifolia* was evaluated using seven primers. The cophenetic correlation of the similarity matrix (r) was 0.7202 indicating that the cluster results had a good fit to the genetic similarity matrix.

Table 2. Total number of bands, number polymorphic bands, percentage of polymorphic bands and polymorphism information content (PIC) generated by 7 primers in 60 accessions of *Elaeagnus latifolia* L. by RAPD analysis

Primer	Primer sequence (5' - 3')	Number of total band	Number of polymorphic band	% PPB	PIC
OPL12	GGG CGG TAC T	29	29	100	0.348
OPO19	GGT GCA CGT T	31	31	100	0.328
OPR04	CCC GTA GCA C	31	30	96.77	0.377
OPT12	GGG TGT GTA G	18	17	94.44	0.267
OPU13	GGC TGG TTC C	27	24	88.88	0.262
OPY04	GGC TGC AAT G	22	19	86.36	0.184
OPAJ18	GGC TGA GTG G	20	18	90.00	0.229
Total		178	168	94.38	0.285




Fig. 2. Banding patterns of 60 accessions of *Elaeagnus latifolia* L. from OPL-12 primer (M: 100 bp marker)




Fig. 3. Banding patterns of 60 accessions of *Elaeagnus latifolia* L. from OPO-19 primer (M: 100 bp marker)




Fig. 4. Dendrogram of 60 accessions of *Elaeagnus latifolia* L. resulting from a UPGMA cluster analysis obtained from seven primers using Jaccard's similarity coefficient.

Discussions

Estimation of genetic diversity among and within crops is crucial in breeding and conservation of genetic resources. RAPD marker analysis is widely used for detection of genetic diversity and classification of the germplasm of many organisms. The different primers are used to amplify different numbers of genotypes. The number of genotypes or accessions amplified by these RAPD primers vary according to the primers used, which depend on the presence of amplification sites (Mahmood *et al.*, 2013). RAPDs have been shown to be an effective method for studying genetic diversity in many plants such as wheat (Al-Rawashdeh, 2011), *Jasminum* species

(Mahmood *et al.*, 2013), mango, *Mangifera indica* (Ramessur and Ranghoo-Sanmukhiya, 2011) and Chilli, *Capsicum annuum* (Peeraulle and Ranghoo-Sanmukhiya, 2013).

In this paper, RAPDs were used to estimate the genetic diversity of *E. latifolia*. Seven primers were selected for characterization of 60 accessions of *E. latifolia*. A total of 178 DNA fragments were identified, 168 were polymorphic. RAPD markers exhibited more than 60 % polymorphism indicating the presence of wide genetic variability among different genotypes and variations in DNA sequences leading to polymorphism. Therefore, this great polymorphism is indicative of great genetic diversity (Gajera *et al.*, 2010). Jaccard's similarity coefficient or genetic similarity (GSj) ranged from 0.38-0.71. The results showed high level of polymorphism occurred in 60 accessions. Fifty eight accessions could be distinguished, while only two accessions; N55 and N72 appeared to be genetically very close with the genetic similarity coefficient of 0.71. Their relatively high similarity may indicate that they developed from a common ancestor with similar in leaf and fruit shapes as well as taste. Clustering patterns of different accessions in the dendrogram revealed that the genetic diversity of *E. latifolia* is not in perfect agreement with geographical diversity. The dendrogram showed that for some groups there was no clear pattern of clustering according to the locations from where accessions were collected. Al-Rawashdeh (2011) reported that genetic similarity was detected by RAPD analysis in the wild medicinal plant *Artemisia judaica*. High similarity indicators of the accessions in the current study suggest that the test entities have a close genetic relation. A slight correlation with geographic location was also noted which is similar to our study for the samples collected from the same location, however the different clusters are identified in the samples collected from the different areas. Same finding was reported by Tonk *et al.* (2011) on *Hypericum perforatum*, clustering based on RAPD data showed a moderate level of correlation with regional origin of the clones.

Conclusion

RAPD markers were found to be useful in the assessment of *E. latifolia* diversity. This understanding of genetic diversity can be used for genetic conservation and genetic resource management of this plant species in the future. Possible paths for future research on genetic diversity in populations of *E. latifolia* is the use of advanced molecular markers techniques such as inter-simple sequence repeats (ISSRs) and/or simple sequence repeats (SSRs).

Acknowledgements

The authors would like to thank Chiang Mai University for financial support and the Science and Technology Research Institute, Chiang Mai University and Plant Biotechnology Research Centre, Royal Project Foundation for providing research facilities.

References

- Ahmadiani, A., Hosseiny, J., Semnani S., Javan, M., Saeedi, F., Kamalinejad, M. and Saremi, S. (2000). Antinociceptive and anti-inflammatory effects of *Elaeagnus angustifolia* fruit extract. *Journal of Ethnopharmacol* 72:287-292.
- Al-Rawashdeh, I. M. (2011). Genetic variability in a medicinal plant *Artemisia judaica* using random amplified polymorphic DNA (RAPD) markers. *International Journal of Agriculture and Biology* 13:279-282.
- Bargali, K. (2011). Actinorhizal plants of Kumaun Himalaya and their ecological significance. *African Journal of Plant Science* 5:401-406.
- Bhattacharyya, B. and Johri, B. M. (1998). Flowering plants taxonomy and phylogeny. Narosa Publishing House, London. pp. 347-349.
- Gajera, B. B., Kumar, N., Singh, A. S., Punvar, B. S., Ravikiran, R., Subhash, N. and Jadeja, G. C. (2010). Assessment of genetic diversity in castor (*Ricinus communis* L.) using RAPD and ISSR markers. *Industrial Crops and Products* 32:491-498.
- Li, H., Ruan, C. J. and Teixeira da Silva, J. A. (2009). Identification and genetic relationship based on ISSR analysis in a germplasm collection of sea buckthorn (*Hippophae* L.) from China and other countries. *Scientia Horticulturae* 123:263-271.
- Mahmood, M. A., Hafiz, I. A., Abbasi, N. A. and Faheem, M. (2013). Detection of genetic diversity in *Jasminum* species through RAPD techniques. *International Journal of Agriculture and Biology* 15:505-510.
- Patel, R. K., Singh, A. and Deka, B. C. (2008). Soh-Shang (*Elaeagnus latifolia*): An under-utilized fruit of North East region needs domestication. *Himalayan Ecology* 16:1-2.
- Peeraullee, N. and Ranghoo-Sanmukhiya, V. M. (2013). Assessment of Genetic Diversity in Local Chilli (*Capsicum annum*) Varieties in Mauritius. *International Journal of Agriculture and Biology* 15:891-896.
- Ramessur, A. D. and Ranghoo-Sanmukhiya, V. M. (2011). RAPD marker-assisted identification of genetic diversity among mango (*Mangifera indica*) varieties in Mauritius. *International Journal of Agriculture and Biology* 13:167-173.
- Rohlf, F. J. (2005). NTSYSpc: Numerical Taxonomy and Multivariate Analysis System, Version 2.2, Exeter Software, Setauket, New York.
- Sakamura, F. and Suga, T. (1987). Changes in chemical components of ripening oleaster fruits. *Phytochemistry* 26:2481-2484.
- Sundriyal, M. and Sundriyal, R. C. (2003). Underutilized edible plants of Sikkim Himalaya: need for domestication. *Current Science* 85:731-736.
- Sundriyal, M. and Sundriyal, R. C. (2005). Seedling growth and survival of selected wild edible fruit species. *Acta Oecologica* 28:11-21.
- Tonk, F. A., Giachino, R. R. A., Sonmez, C., Yuce, S., Bayram, E. M. İ. N. E., Telci, I. and Furan, M. A. (2011). Characterization of various *Hypericum perforatum* clones by hypericin and RAPD analyses. *International Journal of Agriculture and Biology* 13:31-37.

(Received 10 November 2013; accepted 12 January 2014)