
Effect of Chlorine Dioxide (ClO₂) on Culture Medium Sterilization on Micropropagation of Persian Violet (*Exacum affine* Balf.f. ex Regel)

Srichuay, W.^{1*}, Promchan, T.² and Te-chato, S.³

¹Faculty of Science and Technology, Princess of Naradhiwas University, Narathiwat 96000, Thailand; ²Faculty of Agricultural Science and Technology, Silpakorn University, Petchaburee 71260, Thailand; ³Department of Plant Science, Faculty of Natural Resources, Prince of Songkla University, Hatyai, Songkhla 90112, Thailand.

Srichuay, W., Promchan, T. and Te-chato, S. (2018). Effect of chlorine dioxide (ClO₂) on culture medium sterilization on micropropagation of persian violet (*Exacum affine* Balf.f. ex Regel). International Journal of Agricultural Technology 14(2):259-270.

Abstract The effect of chlorine dioxide (ClO₂) on culture medium sterilization in persian violet (*Exacum affine* Balf.f. ex Regel) has been studied. The culture medium was sterilized by ClO₂, which promoted shoot formation. Culture medium supplemented with 5 ppm ClO₂ gave the highest shoot at 9.27±4.13 shoot/explants. The explants were cultured on medium supplemented with 10 ppm ClO₂ which induced the highest root formation at 62.50 % and 3.61±1.00 root/explants significant different with other treatments. In addition to flower induction, above treatment gave the highest flowering at 52.38 % and blooming flower at 14.28 %. In case of concentration of culture medium, ½ MS supplemented with 10 ppm ClO₂ could be root induction at 100 % gave the highest number of root at 12.33±3.84 root/explants significant different with another treatment. The effect of agar concentration in medium, 0.60 % agar gave the best root induction at 94.44 % while 0.75% was induced flower at 52.38 % and blooming flower at 14.28 %. Another study is that effect of ClO₂ when combination with silver nitrate (AgNO₃) in micropropagation. The optimum concentration of AgNO₃ to promoted root induction and flowering was 1.0 mg L⁻¹ at 33.33% root induction, 58.33 % flowering, and 50.00 % blooming flower.

Keywords: Persian violet, chlorine dioxide, sterilization, root induction, flowering

Introduction

Persian violet (*Exacum affine* Balf.f. ex Regel) which belongs to the family *Gentianaceae*, is a widely used indoor or outdoor small potted plant. Persian violet is traditionally propagated by seed and produces fragrant, blue, purple or white coloured flowers (Kapchina-Toteva *et al.*, 2005), but reduced

* **Coressponding author:** Srichuay, W.; **Email:** wutthichai.sri@gmail.com

fertility in cultivars with composed flowers has increased to need for efficient methods of vegetative propagation (Ørnstrup *et al.*, 1993). Normally, *in vitro* *E. affine* Balf culture use flower buds, peduncles (Ørnstrup *et al.*, 1993) and internodal sections (Ballal, 1990) as explants. Many keys are factor affecting successful on tissue culture. Ethylene is important factor in tissue culture and it play both positive and negative role in callus induction, somatic embryogenesis, organogenesis and plant regeneration (Kumar *et al.*, 2009). Ethylene inhibited plant regeneration of elite wheat (Yu *et al.*, 2008). AgNO₃ is one of inhibitor ethylene action (Beyer, 1976) because silver ions might play role in inhibitory effect in ethylene (Fei *et al.*, 2000). Root induction is a finally protocol of tissue culture so root induction is important for survival after transfer to field condition. Genotype and culture medium were also reported to promote root induction (Sanputawong and Te-chato, 2008; Thawaro and Te-chato, 2010; Sirisom and Te-chato, 2014). Gelling agents in culture medium enhanced effect on root formation (Mohan *et al.*, 2005). However, concentration of gelling agents affected to root induction (Suthar *et al.*, 2011).

Normally, eliminate contamination microorganisms from the culture medium is using autoclave at 121 °C and pressure at 1 kg cm⁻²) for 15 to 30 min (Torres *et al.*, 1998). In case of surface explants sterilize before culture, which use mercuric chlorine (Kannan *et al.*, 2007), sodium hypochlorite (Sivanesan *et al.*, 2011), cetrimide (Gantait *et al.*, 2011), and dichloroisocyanuric acid Na₂ salt (Šušek *et al.*, 2002). Chlorine dioxide (ClO₂) is a potent antiprotozoan agent (Peeters *et al.*, 1989), antibacterial agent (Taylor *et al.*, 2000) fungicide (Wilson *et al.*, 2005), and virucide (Taylor and Butler, 1982). Bacteria may be killed effectively by ClO₂ relatively wider range of pH value (pH 3.0-8.0) (Huang *et al.*, 1997). Chlorine ions are generally toxic to plants at relatively low concentrations and may cause irreversible damage to plant development (Carrillo *et al.*, 1996). On the other hand, the benefits of chlorine dioxide in agriculture, which are disinfection of *Escherichia coli* and *Salmonella* sp. on agricultural product surfaces (Han *et al.*, 2000; López-Velasco *et al.*, 2012), postharvest storage quality of plum fruit (*Prunus salicina* L.) (Chen and Zhu, 2011) and sterilization in the *in vitro* culture medium (Cardoso *et al.*, 2012)

Thus, the aim of this study was to evaluate the effects of ClO₂ for chemical sterilization of *in vitro* culture medium and on the development of persian violet plantlets at shoot induction, root induction, and flowering and to compare the growth of these plantlets with plantlets developed using media conventionally sterilized by autoclaving.

Materials and methods

The explants of persian violet (*E. affine* Balf.f. ex Regel) were standard medium [Murashige and Skoog (MS) medium (Murashige and Skoog, 1962) supplemented with 3.0 % (w/v) sucrose, 0.5 mg L⁻¹ BA and 7.5 g l⁻¹ agar, pH 5.8] and subcultured every 1 month. Growth conditions were 28±2° C, under 14 hours of light (23 umolm⁻²s⁻¹ photosynthetic photon flux density).

Effect of chlorine dioxide on shoot induction

The explants were cultured on MS supplemented with 3% sucrose and 0.5 mg L⁻¹ BA and 0.75 % agar, which sterilized by autoclaving (control) and ClO₂ at various concentration (5, 10 and 15 ppm). After 1 month of culture, the percentage of contamination medium and shoot formation, number of shoot/explants and shoot length were recorded. The data were compared with autoclaving and ClO₂ as a sterilizing agent in culture media.

Effect of chlorine dioxide on root and flower induction

The explants were cultured on MS supplemented with 3% sucrose, 0.75 % agar and without BA, which sterilized by autoclaving (control) and various concentration of ClO₂ (5, 10 and 15 ppm). After 1 month of culture, the percentage of contamination medium and root induction, number of root/explants and root length, were recorded. In parameter flower data were collected after 2 month of culture. The data were compared with autoclaving and ClO₂ as a sterilizing agent in culture media.

Effect of concentration agar on culture medium (sterilized by 10 ppm ClO₂) on root and flower induction

The explants were cultured on MS medium supplemented with 3% sucrose, various concentration of agar (0.60, 0.65, 0.70 and 0.75%) and sterilized by 10 ppm ClO₂. After culture, the day of rooted, the percentage of contamination medium, root induction, and number of root/explants and root length were recorded 1 month after incubator and flowering and blooming flower were calculated 2 months after culture.

Effect of concentration of culture medium on root induction

The explants were cultured on different concentration of MS medium (half and full strength concentration), which supplemented with 3% sucrose, 0.75% agar and without BA. All culture medium sterilized by 10 ppm ClO₂. After culture, the day of rooted, the percentage of contamination medium and root induction, and number of root/explants and root length were recorded. The data were compared with autoclaving and ClO₂ as a sterilizing agent in culture media.

Effect of AgNO₃ concentration on culture medium on root and flower induction

The explants were cultured on MS medium supplemented with 3% sucrose, various concentration of AgNO₃ (0, 0.5, 1.0 and 1.5%) solidify with 0.70 % agar and sterilized by 10 ppm ClO₂. After culture, the percentage of contamination medium, root induction, and number of root/explants and root length were recorded 1 month after incubator and flowering and blooming flower were calculated 2 months after culture.

Statistical analysis

All experiments were performed in a completely randomized design (CRD). Each consisted of four replicates per treatment and six explants were performed in each replication. Mean values were analyzed using a one-way analysis of variance (ANOVA). Significant differences among treatments were detected using least significant difference (LSD) with 5 or 1% probability.

Results

Effect of chlorine dioxide on shoot induction

All explants were shoot developed on MS supplemented with 3% sucrose and 0.5 mg L⁻¹ BA (0.75% agar) which sterilized by autoclaving and ClO₂ after culture for 1 month. In case of medium sterilized by ClO₂, medium were not contamination. Shoot could be developed in 5 ppm ClO₂ at 9.27±4.13 shoot/explants. Shoot length was the highest respond on 15 ppm ClO₂ at 7.24±3.77 cm (Table 1).

Effect of chlorine dioxide on root and flower induction

All explants were root developed on MS supplemented with 3% sucrose, 0.75% agar, and sterilized by autoclaving and ClO₂ after culture for 1 month. In

case of medium sterilized by ClO_2 , medium were not contamination. Root could be developed in 10 ppm ClO_2 , which root induction at 62.50% and 3.61 ± 1.00 root/explant. In addition, this concentration gave the highest percentage of flowering and blooming flower at 52.38 and 14.28 %, while 5 ppm ClO_2 gave the highest root length at 7.62 ± 2.44 cm (Table 2).

Effect of concentration of culture medium on root induction

All explants were root response on 1/2MS supplemented with 3% sucrose, 0.70% agar, and sterilized by 10 ppm ClO_2 after culture for 6 days. The best results in percentage root induction at 100%, 12.33 ± 3.84 root/explant, and root length at 19.67 ± 5.10 cm (Table 3) (Figure 1b)

Effect of concentration of agar on root and flower induction

After 1 month of culture, all culture medium cannot contamination. The explants were root response on MS supplemented with 3% sucrose, 0.65% agar, and sterilized by 10 ppm ClO_2 . This culture medium gave the highest root induction at 83.33 %, 6.39 ± 2.23 root/explants, and root length at 7.47 ± 2.01 cm. The term of flowering and blooming flower, explants respond on 0.75 % agar at 52.38 % and 14.28 % (Table 4) (Figure 1c, d and e).

Effect of AgNO_3 concentration on culture medium on root and flower induction

After 1 month of culture, all culture medium cannot contamination. The explants were root induction respond on MS medium supplemented with 3% sucrose, 0.70% agar, 0.5 mg L^{-1} AgNO_3 and sterilized by 10 ppm ClO_2 at 50.00%. While 1.5 mg L^{-1} AgNO_3 gave the highest number of root/explant at 4.21 ± 2.25 and root length at 10.08 ± 5.16 cm. The term of flowering and blooming flower, explants respond on 1.0 mg L^{-1} AgNO_3 at 58.33% and 50.00% (Table 5) (Figure 1g).




Figure 1. *In vitro* propagation of persian violet (*Exacum affine* Balf.f. ex Regel) culture on medium sterilized with 10 ppm ClO_2 (bar : 5 mm). (a) Shoot induction on MS medium with 0.5 BA 0.75% agar. (b) Root formation on PGRs-free $\frac{1}{2}$ MS. Root were induced on PGRs-free MS medium supplemented with 3 different concentration of agar [(c) 0.60%, (d) 0.65% and (e) 0.70% agar]. Blooming flower of complete plantlet culture on PGRs-free MS medium with (f) 0.70% agar and (g) 0.70% agar and $1 \text{ mg L}^{-1} \text{ AgNO}_3$ (h) Complete plantlets developed on PGRs-free MS supplemented with 0.70% agar and $1 \text{ mg L}^{-1} \text{ AgNO}_3$.

Table 1. Effect of different concentration of chlorine dioxide on medium sterilization and shoot formation in persian violet cultured on MS medium supplemented with 3% sucrose and 0.5 mg L⁻¹ BA and 0.75% agar after 1 month of culture. x

TRT	Contamination (%)	Shoot formation (%)	No. of shoot/explant	Shoot length (cm)
Autoclave	0	100	10.67±4.42	9.60±4.08 ^a
5 ppm ClO ₂	0	100	9.27±4.13	5.86±3.03 ^b
10 ppm ClO ₂	0	100	9.07±4.67	3.88±2.01 ^b
15 ppm ClO ₂	0	100	8.40±3.93	7.24±3.77 ^{ab}
F test			ns	**
C.V.(%)			23.15	28.89

Mean values followed by the same letter(s) within a column are not significantly different ($P \leq 0.01$)

Table 2. Effect of different concentration of chlorine dioxide on medium sterilization and root induction in persian violet cultured on PGRs-free MS medium supplemented with 3% sucrose and 0.75% agar after 1-2 month of culture.

TRT	Contamination (%)	Root induction (%)	No. of root/explant	Root length (cm)	Flowering (%)	Blooming flower (%)
Autoclave	0	79.17 ^a	2.60±0.93	3.92±1.60	42.86	0
5 ppm ClO ₂	0	33.33 ^b	3.29±0.58	7.62±2.44	28.57	0
10 ppm ClO ₂	0	62.50 ^b	3.61±1.00	4.93±2.77	52.38	14.28
15 ppm ClO ₂	0	58.33 ^b	2.89±0.81	5.29±3.82	28.57	0
F test		*	ns	ns		
C.V.(%)		65.02	68.86	90.76		

Mean values followed by the same letter(s) within a column are not significantly different ($P \leq 0.01$)

Table 3. Effect of concentration of culture medium on root induction in persian violet cultured on PGRs-free medium supplemented with 3% sucrose and 0.75% agar and sterilized with 10 ppm ClO₂ after 1 month of culture.

TRT	Contamination (%)	Day of rooted	Root induction (%)	No. of root/explant	Root length (cm)
½MS	0	6	100	12.33±3.84 ^a	19.67±5.10 ^a
MS	0	8	83.33	5.42±1.99 ^b	5.37±1.93 ^b
F test			ns	**	**
C.V.(%)			21.51	34.24	30.80

Mean values followed by the same letter(s) within a column are not significantly different ($P \leq 0.01$)

Table 4. Effect of four different concentration of agar on root and flower induction in persian violet cultured on PGRs-free MS medium supplemented with 3% sucrose and 0.75% agar after 1-2 month of culture.

Agar (%)	Contamination (%)	Root induction (%)	No. of root/explant	Root length (cm)	Flowering (%)	Blooming flower (%)
0.60	0	94.44	7.22±3.00 ^a	8.23±3.25 ^a	44.44	0.00
0.65	0	83.33	6.39±2.23 ^{ab}	7.47±2.01 ^a	11.11	0.00
0.70	0	83.33	5.42±1.99 ^{ab}	5.37±1.93 ^{ab}	22.22	8.33
0.75	0	66.67	3.06±1.79 ^b	4.01±3.38 ^b	52.38	14.28
F test		ns	**	*		
C.V.(%)		33.83	41.66	43.53		

Mean values followed by the same letter(s) within a column are not significantly different ($P \leq 0.01$)

Table 5. Effect of different concentration of AgNO₃ on root and flower induction in persian violet cultured on PGRs-free MS medium supplemented with 3% sucrose and 0.75% agar.

AgNO ₃ (mgL ⁻¹)	Contamination (%)	Root induction (%)	No. of root/explant	Root length (cm)	Flowering (%)	Blooming flower (%)
0	0	83.33 ^a	5.42±1.99	5.37±1.93 ^{ab}	22.22	8.33
0.5	0	50.00 ^{ab}	3.96±2.03	3.67±1.96 ^b	33.33	25.00
1.0	0	33.33 ^b	4.00±0.82	4.08±2.18 ^b	58.33	50.00
1.5	0	38.89 ^{ab}	4.21±2.25	10.08±5.16 ^a	25.00	16.67
F test		*	ns	*		
CV.(%)		73.72	60.54	52.04		

Mean values followed by the same letter(s) within a column are not significantly different ($P \leq 0.01$)

Discussion

Shoot formation, in micropropagation of persian violet, ClO₂ touched on shoot induction so, medium sterilized by autoclaving promoted shoot formation then media sterilized with ClO₂. Although ClO₂ increased shoot height, fresh weight and number of leave in gerbera. In case of persian violet, ClO₂ is a few toxic to plant tissue, Srichuay and Te-chato (2014) reported that pineapples were break to growing when they were cultured in medium with high concentration of ClO₂. Effect of ClO₂ on persian violet resulted in sheath blight leaves and shoots were stop developed (Figure don't showed). *In vitro* rooting and flowering are important that plant can adaptation to acclimatization in condition for *in vitro* flowering increased the value of the plant (Yenchon and Te-chato, 2014) and assisted *in vitro* breeding programs (Pratheesh and Kumar, 2012).

Concentration of culture medium on root induction, ½ MS medium promoted a better rooting induction when compared with MS medium. In case of brahmi (*Bacopa monnieri*) half strength MS medium gave the better result in root formation than full strength (Ceasar *et al.*, 2010). ½ MS medium was the best condition for rooted in persian violet. Many researcher reported that many plant species have been induced root formation in half strength MS medium as *Tuberaria major* (Gonçalves *et al.*, 2010), spearmint (Fadel *et al.*, 2010).

Effect of agar on root and flower induction, agar is substance for plant was stood in *in vitro* culture. However concentration of agar has affected to root

induction. High concentration of agar inhibited root development and handicapped root adsorbed water. Suthar *et al.* (2011) reported that high concentration of agar reduced root formation because agar blocked to uptake nutrients in medium.

Concentration of AgNO₃ on root and flower induction, Silver nitrate that break ethylene action inhibited root induction in plant though it promoted flower (Kumar *et al.*, 2009). However, root induction of apple was encouraged by effect of AgNO₃ inhibited ethylene action (Ma *et al.*, 1998). Chithra *et al.* (2004) recorded that AgNO₃ promoted rooting and flowering in rare rheophytic woody medicinal plant whereas rooting of persian violet was interrupted by silver nitrate. AgNO₃ incited flower formation in some plant such as *Capsicum frutescens* (Sharma *et al.*, 2008), *Rosa indica* (Pratheesh and Kumar, 2012).

Acknowledgement

The authors would like to thank National Research Council of Thailand (NRCT), The Center of Excellence in Agricultural and Natural Resources Biotechnology, Department of Plant Science, Faculty of Natural Resources, Prince of Songkla University and Faculty of Science and Technology, Princess of Naradhiwas University for financial support.

References

- Ballal, S. K. (1990). Morphogenic potential of *Exacum affine* and *Eustoma grandiflorum* in tissue culture. *Plant Growth Regulator Society of America Quarterly* 18:73-76.
- Beyer, E. M. (1976). A potent inhibitor of ethylene action in plants. *Plant Physiology* 58:268-271.
- Cardoso, J. C. and Teixeira da Silva, J. A. (2012). Micropropagation of gerbera using chlorine dioxide (ClO₂) to sterilize the culture medium. *In Vitro Cellular & Developmental Biology - Plant* 48:362-368.
- Carrillo, A., Puente, M. E. and Bashan, Y. (1996). Application of diluted chlorine dioxide to radish and lettuce nurseries insignificantly reduced plant development. *Ecotoxicology and Environmental Safety* 35:57-66.
- Cearar, S. A., Maxwell, S. L., Prasad, K. B., Karthigan, M. and Ignacimuthu, S. (2010). Highly efficient shoot regeneration of *Bacopa Monnieri* (L.) using a two-stage culture procedure and assessment of genetic integrity of micropropagated plants by RAPD. *Acta Physiologiae Plantarum* 32:443-52.
- Chen, Z. and Zhu, C. (2011). Combined effects of aqueous chlorine dioxide and ultrasonic treatments on postharvest storage quality of plum fruit (*Prunussalicina* L.). *Postharvest Biology and Technology* 61:117-123.
- Chithra, M., Martin, K. P., Sumandakumari, C. and Madhusoodanan, P. V. (2004). Silver nitrate induced rooting and flowering *in vitro* on rare rheophytic woody medicinal plant, *Rotula aquatica* Lour. *Indian journal of Biotechnology* 3:418-421.
- Fadel, D., Kintzios, S., Economou, A. S., Moschopoulou, G. and Constantinidou, H. I. A. (2010). Effect of different strength of medium on organogenesis, phenolic accumulation

- and antioxidant activity of spearmint (*Mentha spicata* L.) The Open Horticulture Journal 3:31-35.
- Fei, S., Read, P. E. and Riordan, T. P. (2000). Improvement of embryogenic callus induction and shoot regeneration of buffalograss by silver nitrate. *Plant Cell, Tissue and Organ Culture* 60:197-203.
- Gantait, S., Mandal, N., Bhattacharyya, S. and Das, P. K. (2011). Induction and identification of tetraploids using in vitro colchicines treatment of *Gerbera jamesonii* Bolus cv. Sciella. *Plant Cell, Tissue and Organ Culture* 106:485-493.
- Gonçalves, S., Fernandes, L. and Romano, A. (2010). High-Frequency *in Vitro* Propagation of the Endangered Species *Tuberaria Major*. *Plant Cell, Tissue and Organ Culture* 101:359-63.
- Han, Y., Sherman, D. M., Linton, R. H., Nielsen, S. S. and Nelson, P. E. (2000). The effects of washing and chlorinedioxide gas on survival and attachment of *Escherichia coli* O157:H7 to green pepper surfaces. *Food Microbiology* 17:521-533.
- Huang, J., Wang, L., Ren, N., Ma, F. and Ma, J. (1997) Disinfection effect of chlorine dioxide on bacteria in water. *Water Research* 31:607-613.
- Kannan, P., Premkumar, A. and Ignacimuthu. (2007). Thidiazuron induced shoot regeneration in the endangered species, *Exacum travancoricum* Beedi. *Indian Journal of Biotechnology* 6:564-566.
- Kapchina-Toteva, V. M., Iakimova, E. T. and Chavdarov, I. P. (2005). Proceedings of the Balkan Scientific Conference of Biology in Plovdiv (Bulgaria). 19-21 May 2005. pp. 714-722.
- Kumar, V., Parvatam, G. and Ravishankar, G. A. (2009). AgNO₃-a potential regulator of ethylene activity and plant growth modulator. *Journal of Biotechnology* 2:1-15.
- López-Velasco, G., Tomás-Calleja, A., Sbodio, A., Artés-Hernández, F. and Suslow, T. V. (2012). Chlorine dioxide dose, water quality and temperature affect the oxidative status of tomato processing water and its ability to inactivate *Salmonella*. *Food Control* 26: 28-35.
- Ma, J. H., Yao, J. L., Cohen, D. and Morris, B. (1998). Ethylene inhibitors enhance in vitro root formation from apple shoot cultures. *Plant Cell Reports* 17:211-214.
- Murashige, T. and Skoog, F. (1962). A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum* 15:473-497.
- Mohan, R., Chui, E. A., Biasi, L. A. and Soccol, C. R. (2005). Alternative *in vitro* propagation: use of sugarcane bagasse as a low cost support material during rooting stage of strawberry Cv. Dover. *Brazilian Archives of Biology and Technology* 48:37-42.
- Ørnstrup, H., Mølgaard, J. P. and Farestveit, H. (1993), Somatic embryogenesis and plant regeneration from cell suspensions of *Exacum affine*. *Plant Cell, Tissue and Organ Culture* 35:37-41.
- Peeters, J. E., Mazás, A. E., Masschelein, W. J., de Maturana, V. M. and Debacker, E. (1989) Effect of disinfection of drinking water with ozone or chlorine dioxide on survival of *Cryptosporidium parvum* oocysts. *Applied and Environmental Microbiology* 55:1519-1522.
- Pratheesh, P. T. and Kumar, M. A. (2012). *In vitro* flowering in *Rosa indica* L. *International Journal of Pharmacy and Biological Science* 2:196-200.
- Sanputawong, S. and Te-chato, S. (2008). Effect of genotypes of oil palm as indicator for speed of callus and embryogenic callus formation. *Journal of Agricultural Technology* 4:147-156.

- Sharma, A., Kumar, V., Giridhar, P. and Ravishankar, G. A. (2008). Induction of *in Vitro* Flowering in *Capsicum frutescens* under the Influence of Silver Nitrate and Cobalt Chloride and Pollen Transformation. *Electronic Journal of Biotechnology* 11:1-6.
- Sivanesan, I., Song, J. Y., Hwang, S. J. and Jeong, B. R. (2011). Micropropagation of *Cotoneaster wilsonii* Nakai—a rare endemic ornamental plant. *Plant Cell, Tissue and Organ Culture* 105:55-63.
- Sirisom, Y. and Te-chato, S. (2014). Assessment of somaclonal variations of *in vitro*-plants derived from nodal culture of rubber trees by SSR markers. *Songklanakarin Journal of Plant Science* 1:7-12.
- Srichuay, W. and Te-chato, S. (2014). Effect of chlorine dioxide (ClO₂) on sterilization in micropropagation of pineapple cv. Phulae by bioreactor system. *Khon Kaen Agriculture Journal* 42(Suppl 3):75-80.
- Šušek, A., Javornik, B. and Bohanec, B. (2002). Factors affecting direct organogenesis from flower explants of *Allium giganteum*. *Plant Cell, Tissue and Organ Culture* 68:27-33.
- Suthar, R. K., Habibi, N. and Purohit, S. D. (2011). Influence of agar concentration and liquid medium on *in vitro* propagation of *Boswellia serrata* Roxb. *Indian Journal of Biotechnology* 10:224-227.
- Taylor, G. R. and Butler, M. (1982). A comparison of the virucidal properties of chlorine, chlorine dioxide, bromine chloride and iodine. *Journal of Hygiene* 89:321-328.
- Taylor, R. H., Falkinham, J. O., Norton, C. D. and LeChevallier, M. W. (2000). Chlorine, chloramine, chlorine dioxide, and ozone susceptibility of *Mycobacterium avium*. *Applied and Environmental Microbiology* 66:1702-1705.
- Thawaro, S. and Te-chato, S. (2010). Effect of culture medium and genotype on germination of hybrid oil palm zygotic embryos. *ScienceAsia* 36:26-32.
- Torres, A. C., Caldas, L. S. and Buso, J. A. (1998). *Cultura de tecidos e transformação genética de plantas (Tissue culture and genetic transformation of plants)*. EMBRAPA-SPI and CNPH, Brasilia.
- Wilson, S. C., Wu, C., Andriychuk, L. A., Martin, J. M., Brasel, T. L., Jumper, C. A. and Straus, D. C. (2005). Effect of chlorine dioxide gas on fungi and mycotoxins associated with sick building syndrome. *Applied Environment Microbiology* 71:5399-5403.
- Yenchon, S. and S. Te-chato. (2014). Enhanced efficiency of flowering of Rose (*Rosa hybrida*) cv. “My Valentine” *in vitro*. *Songklanakarin Journal of Plant Science* 1:31.34.
- Yuan, Y. and Xu, D. Q. (2001). Stimulation effect of gibberellic acid short-term treatment on leaf photosynthesis related to the increase in Rubisco content in board bean and soybean. *Photosynthesis Research* 68:39-47.

(Received: 5 September 2017, accepted: 20 February 2018)