

---

## Effect of *Moringa oleifera* Seed Oil, Root and Leaf Extracts on Growth of Major Pathogenic Fungi of Tomato, Green Bean and Potato in Vitro

---

El-Mohamedy, R. S. R.\* and Mohamed, S. K.

Plant Pathology Department, National Research Center, Dokki, Cairo, Egypt.

El-Mohamedy, R. S. R. and Mohamed, S. K. (2018). Effect of *Moringa oleifera* seed oil, root and leaf extracts on growth of major pathogenic fungi of tomato, green bean and potato in vitro. International Journal of Agricultural Technology 14(4):505-520.

**Abstract** Fungal plant diseases cause great losses in yield and products of the important vegetable crops in Egypt. In this study, the antifungal activity of *Moringa oleifera* seed oil (MSO), roots extract (MRE) and leaves extract (MLE) against linear growth, spores and sclerotia germination of 18 pathogenic fungi incited tomato, green bean and potato plants was investigated. Crude extracts of roots, leaves as well as seed oil of *Moringa oleifera* significantly reduced radial growth, spores and sclerotia germination of all the tested fungi. *M. oleifera* seed oil and their extracts had different degrees of reduction in both growth and sporulation of the tested fungi. The complete reduction of radial growth, spores and sclerotia germination were at high concentrations of MLE (50%), MRE (20%) and MSO (3%) with all tested fungi. *Fusarium oxysporum*, *F. solani* and *Alternaria solani* were highly affected by MRE, MLE and MSO than *Rhizoctonia solani*, *Sclerotium rolfsii*, *Sclerotinia sclerotium* and *M. phaseolina*. It could be suggested that MRE, MLE and MSO of *Moringa oleifera* could be easily applied as a natural fungicide against fungal pathogens of many important crops.

**Keywords:** *Moringa oleifera*, phytopathogenic fungi, Antifungal activity, plant extract

### Introduction

Fungal infections cause significant loss in many economic crops (Agrios 2005). Many plant pathogenic fungi incited tomato, potato and green bean, such pathogens cause great losses in yield and products during vegetative growth or after harvesting. Chemical control may be available to effectively and extensively reduce the negative effects of such fungal diseases, but field application of these chemical fungicides may not always be desirable. Excessive and improper use of these fungicides presents a danger to the health of humans, animals, and the environment. Therefore, extensive searches for natural and bio fungicides that are environmentally safe and easily biodegradable have been carried out during the last two decades

---

\* **Corresponding Author:** El-Mohamedy, R. S. R.; **E-mail:** riadelmohamedy@yahoo.com

(Gnanamanickam, 2002, Zaker, 2016; Seem *et al.*, 2011; El-Mohamedy *et al.*, 2017). Various plant products like plant extracts, essential oils, gums, resins *etc.* were shown to exert biological activity *in vitro* and *in vivo* and are used as bio-fungicidal compounds, (Fawzi *et al.*, 2009; Al-Askar and Rashad, 2010). Many essential oils and plant extracts have been reported to have antifungal activities with no side effects on humans and animals (Rai and Carpinella, 2006; Tabassum and Vidyasagar, 2013). Essential oils may have a minimum adverse effect on the physiological processes of plants and have less environmental hazards compared to their synthetic alternatives. Since essential oils are plant products, they are easily convertible into a common organic material and eco-friendly (Gnanamanickam, 2002; Seema *et al.*, 2011).

Moringa (*Moringa oleifera* Lam.) has gained much importance in the recent days due to its multiple used and benefits to agriculture and industry. Regarded as a miracle plant, all parts of moringa plant are used for medicinal and other purposes (Price, 2000; Anwar *et al.*, 2007; Garima *et al.*, 2011). In recent years, interests have been generated in the development of safer antifungal agents from natural plant products such as, essential oils and extracts to control fungal plant diseases. The fungicidal effect of Moringa extracts on some soil-borne fungi such as *Rhizoctonia*, *Pythium* and *Fusarium* was recorded by many investigators (Moyo *et al.*, 2012). Dwivedi and Enespa (2012) indicate that *Moringa oleifera* extracts (leaves, bark and seeds) 75 % (v/v) showed significant inhibition in the mycelial growth of *Fusarium solani* and *Fusarium oxysporum* f. sp. *Lycopersici*. *Moringa oleifera* provides a rich and rare combination of zeatin, quercetin, b-sitsterol, caffeoylquinic acid and kaempferol which have antifungal and antibacterial activities (Anjorin *et al.*, 2010). Moringa leaf extract is best used as plant growth enhancer, insect repellent and fungicide (Phiri and Mbewe, 2010)

The aim of this work was to investigate the antifungal activity of *Moringa oleifera* seed oil (MSO), leaves extract (MLE) and roots extract (MRE) against 18 phytopathogenic fungi the causal agents of many fungal diseases tomato, green bean and potato *in vitro*.

## **Materials and methods**

### ***Plant materials***

This study was carried out at the Department of Plant Pathology, National Research Center, Egypt. The source of *Moringa oleifera* seed oil (MSO), leaves extract (MLE) and roots extract (MRE) were kindly prepared and obtained

from Egyptian Scientific Society of Moringa (ESSM), National Research Center, Dokki, and Cairo, Egypt.

### ***Pathogenic fungi***

The 18 isolates of pathogenic fungi were investigated which belonging to eight fungal genera i.e., *Fusarium*, *Rhizoctonia*, *Alternaria*, *Phytophthora*, *Sclerotium*, *Sclerotinia*, *Botrytis* and *Macrophomina* representing 11 species as follows : Seven isolates of tomato pathogenic fungi i.e., *Fusarium oxysporum* f. sp. *lycopersici* (Fol2), *F. oxysporum* f. *radicis* (For15), *F. solani* (Fs 5), *Rhizoctonia solani* (Rs 5), *Sclerotium rolfsii* (Sr 5), *Alternaria solani* (As 3) and *Phytophthora infestans* (Ph 3) & Seven isolates of pathogenic fungi of green bean i.e., *Fusarium oxysporum* (Fox 1), *Fusarium solani* (Fs 5), *Rhizoctonia solani* (Rs 5), *Macrophomina phaseolin* (Mph 3), *Sclerotium rolfsii* (Sr 5), *Botrytis cinerea*, *Sclerotinia sclerotiorum* as well as four isolates of potato pathogenic fungi i.e., *Fusarium semibaticum*, *Rhizoctonia solani* (Rs 5), *Phytophthora infestans* and *Alternaria solani* were maintained and grown on potato dextrose agar medium. These fungal isolates were previously isolated and identified at Plant Pathology Department, National Research Center, Cairo, Egypt. The pathogenicity of each fungi was tested and recorded in previous studies by the author (El-Mohamedy *et al.*, 2013).

### ***Antifungal activity assay***

Antifungal activity of *Moringa oleifera* seed oil (MSO), leaves extract (MLE) and roots extract (MRE) against the tested plant pathogenic fungi mentioned above was carried out using poison food technique (Nene and Thaplyal, 1979) and performed by the agar medium assay. Potato dextrose agar (PDA) medium with different concentrations of moringa leaf extracts (MLE) i.e., 10, 20, 30, 40, 50 % and roots extract (MRE) i.e., 5,10,15,20 and 25% as well as different concentrations of moringa seed oil (MSO) i.e., 1.0, 1.5, 2.0, 2.5 and 3.0 % were prepared by adding appropriate quantity of moringa seed oil to melted medium, followed by addition of Tween 80 (100  $\mu$ L to 100 mL of medium) to disperse the oil in the medium. About 20 ml of the medium were poured into glass Petri-dishes (9 cm x 1.5 cm). A 5 mm diameter agar disk bearing hyphae 7-days-old colonies grown on PDA medium of each tested pathogenic fungi was transferred at the center of each Petri-dish. Positive control (without treatments) plates were inoculated following the same procedure. Plates were incubated at 25  $\pm$  2  $^{\circ}$ C for 8 days and the colony diameter was recorded each day. The MGI (Mycelia Growth Inhibition) percentage was

calculated and expressed as percentage of reduction. Sclerotia germination of *R. solani*, *S. rolfsii*, *M. phaseolina*, *Botrytis cinerea* and *Sclerotinia sclerotiorum* produced on potato dextrose agar (PDA) in each treatment as maintained above were determined according to Manning *et al.* (1970).

Meanwhile, For *F. oxysporum*, *F. solani*, *F. semibaticum*, *A. solani* and *Phytophthora infestans* microscope slides were covered with 1 mL of spore's suspension of each tested pathogen in aqueous solution of the desired moringa extracts or moringai seed oil concentrations in Petri dishes, then incubated at  $27 \pm 1$  °C for 8 h in complete darkness. The percentage of germination was assessed according to El-Abyad and Saleh (1917). Five plates were prepared for each treatment and the means were compared. Antifungal activity was calculated and expressed as percentage of reduction in both linear growth (LG) as well as on spores and sclerotia germination (SG and SCG) of each pathogenic fungi under investigations.

### ***Statistical analyses***

The data of this study were subjected to analysis of variance using the statistical analysis software (Co Stat, 2005). Comparisons among means were made using Duncan's multiple range test (Duncan, 1955).

## **Results**

The antifungal activity of Moringa leaves extract (MLE) at five concentrations i.e., 10, 20, 30, 40 and 50 % (w/v), Moringa roots extract (MRE) at 5, 10, 15, 20 and 25 % (w/v) as well as *Moringa oleifera* seed oil (MSO) at 1.0, 1.5, 2.0, 2.0 and 3.0 % (w/v) concentrations against linear growth (LG), spores and sclerotia germination (SG and SCG) of 18 pathogenic fungal isolates, the causal agents of major fungal disease on tomato, green bean and potato crops was evaluated *in vitro*.

### ***The antifungal activity of Moringa leaves extract (MLE)***

The inhibitory effect of MLE at five concentrations i.e., 10, 20, 30, 40 and 50 % (w/v) against linear growth (LG) as well as spores and sclerotia germination (SG and SCG) of 18 isolates of pathogenic fungi of tomato, green bean and potato crops was investigated and the linear growth and their reduction as well as reduction in spores and sclerotia germination were illustrated in Tables (1 and 2).

**Table 1.** Reduction % in linear growth (LG) of pathogenic fungi on tomato, green bean and potato as affected by different concentrations of moringa leaves extract ( MLE ) on PDA medium

Isolate of Pathogenic fungi	Host plant	MLE concentration									
		10%		20%		30%		40%		50%	
		LG	R	LG	R	LG	R	LG	R	LG	R
<i>Fusarium ox f. sp. lycopersici</i> (Fol2)	Tomato	63.0	30.0	52.0c	42.2	36.0c	60.0	19.8d	78.0	0.0b	100
<i>F. ox radialis. lycopersici</i> (For15)	Tomato	59.0	34.4	46.9d	47.8	34.0c	62.2	19.8d	78.0	0.0b	100
<i>Fusarium oxysporum</i> (Fox 1)	Green bean	53.0	41.1	46.2d	48.6	28.6d	68.2	0.0e	100	0.0b	100
<i>F. solani</i> (Fs 5)	Tomato	58.1	35.5	52.0c	42.2	34.0c	62.2	18.0d	80.0	0.0b	100
<i>F. solani</i> (Fs 3)	Green bean	50.0	44.4	36.0e	60.0	20.7d	77.0	0.0e	100	0.0b	100
<i>Fusarium semibaticum</i> (Fse 2)	Potato	58.1	35.5	46.8d	48.0	34.3c	61.8	25.0c	72.2	0.0b	100
<i>Alternaria solani</i> (As 3)	Tomato	61.0	32.2	54.0b	40.0	42.3b	53.0	26.8c	70.2	0.0b	100
<i>Alternaria solani</i> (As 1)	Potato	63.0	30.0	55.4b	38.4	39.7b	55.8	30.6b	66.0	0.0b	100
<i>Phytophthora infestans</i> (Ph 3)	Tomato	64.3	28.5	53.2b	40.8	41.4b	54.0	26.6c	70.4	0.0b	100
<i>Phytophthora infestans</i> (Ph 1)	Potato	62.1	31.0	50.4c	44.0	38.5c	57.2	21.6c	76.0	0.0b	100
<i>Rhizoctonia solani</i> (Rs 5)	Tomato	67.3	25.2	54.0c	40.0	44.6b	50.4	32.4b	64.0	0.0b	100
<i>Rhizoctonia solani</i> (Rs 1)	Green bean	63.0	30.0	50.0c	44.4	42.4b	52.8	30.2b	66.4	0.0b	100
<i>Rhizoctonia solani</i> (Rs 2)	Potato	65.7	27.0	54.0c	40.0	39.6b	56.0	27.0c	70.0	0.0b	100
<i>Sclerotium rolfsii</i> (Sr 5)	Potato	69.7	22.5	59.4b	34.4	43.3b	51.8	36.0b	60.0	0.0b	100
<i>Sclerotium rolfsii</i> (Sr 3)	Green bean	71.7	20.8	66.6b	26.0	42.4b	52.8	30.4b	66.2	0.0b	100
<i>Sclerotinia sclerotiorum</i> (Sc 2)	Green bean	70.2	22.0	57.6b	36.0	44.8b	50.2	32.0b	64.4	0.0b	100
<i>Botrytis cinerea</i> (Bc 3)	Green bean	64.6	28.2	53.2c	40.8	41.2b	54.2	28.8c	68.0	0.0b	100
<i>Macrophomina phaseolina</i> (Mph 3)	Green bean	61.0	32.2	50.0c	44.4	37.8	58.0	20.7d	77.0	0.0b	100
Control		90.0a	0.0	90.0a	0.0	90.0a	0.0	90.0a	0.0	90.0a	0.0

LG = colony diameter (mm). R = Inhibition percentage (%) as compared to the control. Means followed by the same letters are not significantly different according to Duncan's multiple range test ( $P \leq 0.05$ ).

**Table 2.** Reduction % in spores and sclerotia germination of Pathogenic fungi on tomato, green bean and potato as affected by different concentration of MLE on PDA medium

Isolate of Pathogenic fungi	Host plant	MLE concentration				
		10%	20%	30%	40%	50%
Spores germination						
<i>Fusarium ox</i> f. sp. <i>lycopersici</i> (Fol2)	Tomato	33.8	47.6	66.8	80.2	100
<i>F. ox radialis. lycopersici</i> (For15)	Tomato	38.0	42.6	67.0	81.8	100
<i>Fusarium oxysporium</i> (Fox 1)	Green bean	42.4	48.6	78.0	100	100
<i>F. solani</i> (Fs 5)	Tomato	38.2	48.0	70.0	90.0	100
<i>F. solani</i> (Fs 3)	Green bean	48.0	53.2	78.2	100	100
<i>Fusarium semibaticum</i> (Fse 2)	Potato	38.2	50.4	68.0	100	100
<i>Alternaria solani</i> (As 3)	Tomato	34.8	44.8	58.2	74.2	100
<i>Alternaria solani</i> (As 1)	Potato	32.2	40.0	52.4	70.8	100
<i>Phytophthora infestans</i> (Ph 3)	Tomato	30.2	44.2	60.0	74.4	100
<i>Phytophthora infestans</i> (Ph 1)	Potato	33.8	50.2	64.2	77.4	100
Sclerotia germination						
<i>Rhizoctonia solani</i> (Rs 5)	Tomato	21.8	30.8	48.0	60.0	100
<i>Rhizoctonia solani</i> (Rs 1)	Green bean	24.6	32.0	46.2	62.0	100
<i>Rhizoctonia solani</i> (Rs 2)	Potato	24.4	36.0	51.0	66.4	100
<i>Sclerotium rolfsii</i> (Sr 5)	Potato	21.0	27.0	42.2	60.8	100
<i>Sclerotium rolfsii</i> (Sr 3)	Green bean	18.4	25.0	38.8	58.4	100
<i>Sclerotinia sclerotiorum</i> (Sc 2)	Green bean	20.0	32.0	44.4	61.2	100
<i>Botrytis cinerea</i> (Bc 3)	Green bean	23.4	34.2	48.0	60.0	100
<i>Macrophomina phaseolina</i> (Mph 3)		30.8	42.4	58.0	74.8	100

All tested concentrations of MLE caused significant decrease in linear growth (mm) of all tested pathogenic fungi. The inhibitory of MLE increased by increasing its concentrations to reach to 100% reduction at 50 % concentration is demonstrated in Table 1. MLE at 40% concentration inhibited the linear growth by ranging from 100 – 60.0 %, however, at 30 % it causes reduction ranging from 77.2 -51.8%. Meanwhile MLE at 20 % and 10 % the least antifungal activity (<50% reduction) of all tested fungi. The highest records in linear growth reduction were with all isolates of *Fusarium*, *Phytophthora*, *Alternaria* followed by *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium* in ascending order.

Results clearly showed that all tested concentrations of MLE had the ability to decrease the germination of both spores and sclerotia of the tested fungi with different values (Table 2). MLE at 50 % concentration caused complete reduction in spores and sclerotia germination at 40% concentration caused 100-70% and 74.8-58.4 % respectively in reduction of both spores and sclerotia germination. Meanwhile, at 20 % of MLE the moderate reduction ( up to 53 of spores and up to 42.2% of sclerotia germination were recorded. The

least effect (< 50%) was at 10 % of MLE. Spores of all tested isolates of *Fusarium*, *Phytophthora* and *Alternaria* were highly affected with MLE than the sclerotia of *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium*. It was observed that all tested isolates of *Fusarium*, *Phytophthora* and *Alternaria* were more sensitive to all concentrations of MLE than *Macrophomina* and *Rhizoctonia*, meanwhile, isolates of *Botrytis*, *Sclerotinia* and *Sclerotium* were least affected by MLE (Tables 1 and 2).

### ***The antifungal activity of Moringa root extract (MRE)***

The inhibitory effect of MRE at five concentrations i.e., 5, 10, 15, 20 and 20 % (w/v) on linear growth (LG) as well as spores and sclerotia germination (SG and SCG) of 18 isolates of pathogenic fungi of tomato, green bean and potato crops was investigated and the linear growth and their reduction as well as reduction in spores and sclerotia germination were illustrated in Tables 3 and 4.

Results demonstrated that MRE at all tested concentrations showed significantly inhibitory effects against the linear growth (LG) of all tested pathogenic fungi. The reduction in linear growth was increased by increasing the concentration of MRE (Table 3). All fungal isolates of *Fusarium*, *Phytophthora* and *Alternaria* were highly affected with all concentrations of MRE more than the other tested fungi. MRE at 25 % caused complete reduction of the linear growth (LG) of all tested fungi, however, at 20% the reduction percentages ranging from 100 – 79.0 % with *Fusarium*, *Phytophthora* and *Alternaria*, and by 80.0-66.0% with *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium*. Meanwhile, at 15% concentration of MRE the reduction in linear growth of such fungi ranging from 85.0—62.0% and 71.8-48.0% respectively were recorded. MRE at 5% showed the least records of linear growth reduction with all tested fungi.

All tested concentrations of MLE could decrease germination of spores and/or sclerotia of the tested pathogenic fungi (Table 4). This decreasing in spores and/or sclerotia germination reached to 100% reduction of all tested fungi at 25% concentration of MRE, however at 20% concentration the reduction ranging 100-88.2 % and 90.0 - 65.2% of spores and sclerotia germination. Meanwhile MRE at 15 % caused reduction ranging from 94.4 - 70.6% and 82.2-60.0 % of spores and sclerotia germination respectively. However, the moderate effects on spores and sclerotia germination (78.4-56.6% and 66.4-37.8% reduction) were recorded at 10 % of MRE. The least effects on the germination of each spores and sclerotia of the tested fungi were at 5 % of MRE. Spores of *Fusarium*, *Phytophthora* and *Alternaria* were more affected

with MRE than sclerotia of *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium*.

### ***The antifungal activity of Moringa seed oil (MSO)***

The inhibitory effect of MSO at five concentrations i.e., 1.0, 1.5, 2.0, 2.5 and 3.0 % (w/v) on linear growth (LG) as well as spores and sclerotia germination (SG and SCG) of 18 isolates of pathogenic fungi of tomato, green bean and potato crops was investigated and the linear growth and their reduction as well as reduction in spores and sclerotia germination were illustrated in Tables 5 and 6. Results showed that all tested concentrations of MSO were significantly inhibited the linear growth (LG) of all tested pathogenic fungi with different degrees of reduction. MSO at 3.0 % completely reduction in the linear growth (LG) of all tested fungi were recorded, but at 2.5 % the reduction reached to 100% of all isolates of *Fusarium*, *Phytophthora* and *Alternaria*, but ranging from 88.0 - 62.0% of *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium*. Meanwhile, MSO at 2.0 % caused linear growth reduction of all of *Fusarium*, *Phytophthora* and *Alternaria* tested isolates ranging from 84.2-62.2 % and by 62.0 - 42.0 % of *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium* were noted. However, the reduction reached 72.0 - 50.8% and 56.0 - 40.2% of the same pathogenic fungal isolates respectively at 1.5 % concentration.

Results demonstrated that MSO at all tested concentrations that resulted in decreasing spores and sclerotia germination of all tested fungi to reach to 100 % reduction at 3.0 % concentration of MSO (Table 6). Spores of *Fusarium*, *Phytophthora* and *Alternaria* was highly affected by all tested MSO concentration when compared with sclerotia of *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium*. Meanwhile, at 2.5 % of MSO the reduction reached to 100% of spore germination with *Fusarium*, *Phytophthora* and *Alternaria*, but ranging from 74.0 – 64.2 % of *Macrophomina*, *Rhizoctonia*, *Botrytis*, *Sclerotinia* and *Sclerotium* that at 2.0% concentration the reduction reached to 90.0-68.2% and 70.0-55.0% of the same pathogenic fungi respectively. The moderate effect of MSO was at 1.5 % concentration. It decreased the spores and sclerotia germination of all tested fungi to reach 77.8-58.0% and 57.4 - 50.8% respectively. However, the reduction of spores and sclerotia germination ranging from 60.2-52.2% and 48.2-32.8% at 1.0% of MSO with all tested fungi were noted.

**Table 3.** Reduction % in linear growth (LG) of plant pathogenic fungi as affected by different concentrations of moringa leaves extract (MRE) on PDA medium

Isolate of Pathogenic fungi	Host plant	MRE concentration									
		5%		10%		15%		20%		25%	
		LG	R	LG	R	LG	R	LG	R	LG	R
<i>Fusarium ox</i> f. sp. <i>lycopersici</i> (Fol2)	Tomato	52.4b	41.7	36.0d	60.0	27.0d	70.0	13.5c	85.0	0.0b	100
<i>F. ox radialis. lycopersici</i> (For15)	Tomato	50.4b	44.0	36.0d	60.0	23.2d	74.2	10.8c	88.0	0.0b	100
<i>Fusarium oxysporum</i> (Fox 1)	Green bean	52.1b	42.1	36.0d	60.6	23.1d	74.3	9.0d	90.0	0.0b	100
<i>F. solani</i> (Fs 5)	Tomato	46.8d	48.0	30.9e	65.6	17.1e	81.0	0.0d	100	0.0b	100
<i>F. solani</i> (Fs 3)	Green bean	49.0c	45.5	27.0e	70.0	13.5e	85.0	0.0d	100	0.0b	100
<i>Fusarium semibaticum</i> (Fse 2)	Potato	52.4b	41.7	36.0d	60.0	22.5d	75.0	5.4d	94.0	0.0b	100
<i>Alternaria solani</i> (As 3)	Tomato	50.0c	45.0	43.2c	52.4	27.0d	70.0	14.4c	84.0	0.0b	100
<i>Alternaria solani</i> (As 1)	Potato	55.8c	38.0	45.0c	50.0	34.2c	62.0	17.2c	80.8	0.0b	100
<i>Phytophthora infestans</i> (Ph 3)	Tomato	53.4c	40.6	39.6d	56.0	26.8d	70.2	27.0b	79.0	0.0b	100
<i>Phytophthora infestans</i> (Ph 1)	Potato	50.9c	43.4	35.2d	60.8	19.1e	78.8	14.4c	84.0	0.0b	100
<i>Rhizoctonia solani</i> (Rs 5)	Tomato	56.9c	36.7	43.2c	52.0	32.4c	64.0	25.2b	72.0	0.0b	100
<i>Rhizoctonia solani</i> (Rs 1)	Green bean	58.5b	35.0	45.0c	50.0	28.9dc	67.8	22.5b	75.0	0.0b	100
<i>Rhizoctonia solani</i> (Rs 2)	Potato	55.8c	38.0	40.5c	55.0	25.0d	72.2	18.0c	80.0	0.0b	100
<i>Sclerotium rolfsii</i> (Sr 5)	Potato	63.0b	30.0	50.0b	44.4	47.6b	58.2	27.0b	70.0	0.0b	100
<i>Sclerotium rolfsii</i> (Sr 3)	Green bean	64.9b	27.8	53.0b	30.0	46.8b	48.0	30.6b	66.0	0.0b	100
<i>Sclerotina sclerotiorum</i> (Sc 2)	Green bean	58.5b	35.0	46.6b	48.2	38.8c	56.8	18.0	80.0	0.0b	100
<i>Botrytis cinerea</i> (Bc 3)	Green bean	57.4b	36.2	46.0b	48.8	34.2c	62.2	23.4b	74.0	0.0b	100
<i>Macrophomina phaseolina</i> (Mph 3)	Green bean	53.8c	40.2	37.4	58.4	25.3d	71.8	18.0c	80.0	0.0b	100
Control		90.0a	0.0	90.0a	0.0	90.0a	0.0	90.0a	0.0	90.0a	0.0

LG = colony diameter (mm) . R = Inhibition percentage (%) as compared to the control. Means followed by the same letters are not significantly different according to Duncan's multiple range test ( $P \leq 0.05$ ).

**Table 4.** Reduction % in spores and sclerotia germination of pathogenic fungi on tomato, green bean and potato as affected by different concentration of MRE on PDA medium

Isolate of Pathogenic fungi	Host plant	MRE concentration				
		5%	10%	15%	20%	25%
Spores germination						
<i>Fusarium ox f. sp. lycopersici</i> (Fol2)	<b>Tomato</b>	<b>50.4</b>	<b>67.2</b>	<b>82.0</b>	<b>92.2</b>	<b>100</b>
<i>F. ox radicans. lycopersici</i> (For15)	<b>Tomato</b>	<b>52.2</b>	<b>68.2</b>	<b>88.2</b>	<b>90.0</b>	<b>100</b>
<i>Fusarium oxysporium</i> (Fox 1)	<b>Green bean</b>	<b>50.4</b>	<b>73.2</b>	<b>92.8</b>	<b>100</b>	<b>100</b>
<i>F. solani</i> (Fs 5)	<b>Tomato</b>	<b>58.8</b>	<b>74.8</b>	<b>90.0</b>	<b>100</b>	<b>100</b>
<i>F. solani</i> (Fs 3)	<b>Green bean</b>	<b>60.4</b>	<b>78.4</b>	<b>94.0</b>	<b>100</b>	<b>100</b>
<i>Fusarium semibaticum</i> (Fse 2)	<b>Potato</b>	<b>50.2</b>	<b>68.2</b>	<b>84.2</b>	<b>100</b>	<b>100</b>
<i>Alternaria solani</i> (As 3)	<b>Tomato</b>	<b>50.6</b>	<b>58.4</b>	<b>80.0</b>	<b>90.0</b>	<b>100</b>
<i>Alternaria solani</i> (As 1)	<b>Potato</b>	<b>44.8</b>	<b>56.6</b>	<b>70.0</b>	<b>88.2</b>	<b>100</b>
<i>Phytophthora infestans</i> (Ph 3)	<b>Tomato</b>	<b>46.0</b>	<b>60.4</b>	<b>77.0</b>	<b>88.0</b>	<b>100</b>
<i>Phytophthora infestans</i> (Ph 1)	<b>Potato</b>	<b>52.6</b>	<b>64.0</b>	<b>80.0</b>	<b>92.2</b>	<b>100</b>
Sclerotia germination						
<i>Rhizoctonia solani</i> (Rs 5)	<b>Tomato</b>	<b>40.2</b>	<b>58.2</b>	<b>70.0</b>	<b>73.0</b>	<b>100</b>
<i>Rhizoctonia solani</i> (Rs 1)	<b>Green bean</b>	<b>42.0</b>	<b>53.8</b>	<b>68.2</b>	<b>80.0</b>	<b>100</b>
<i>Rhizoctonia solani</i> (Rs 2)	<b>Potato</b>	<b>46.2</b>	<b>58.4</b>	<b>80.2</b>	<b>84.4</b>	<b>100</b>
<i>Sclerotium rolfsii</i> (Sr 5)	<b>Potato</b>	<b>32.7</b>	<b>40.2</b>	<b>50.2</b>	<b>66.8</b>	<b>100</b>
<i>Sclerotium rolfsii</i> (Sr 3)	<b>Green bean</b>	<b>30.4</b>	<b>37.8</b>	<b>60.0</b>	<b>65.2</b>	<b>100</b>
<i>Sclerotina sclerotiorum</i> (Sc 2)	<b>Green bean</b>	<b>33.6</b>	<b>52.8</b>	<b>74.0</b>	<b>80.0</b>	<b>100</b>
<i>Botrytis cinerea</i> (Bc 3)	<b>Green bean</b>	<b>38.8</b>	<b>50.8</b>	<b>65.4</b>	<b>78.0</b>	<b>100</b>
<i>Macrophomina phaseolina</i> (Mph 3)		<b>48.8</b>	<b>66.4</b>	<b>82.4</b>	<b>90.0</b>	<b>100</b>

**Table 5.** Reduction % in linear growth (LG) of plant pathogenic fungi on tomato, green bean and potato as affected by different concentrations of moringa leaves extract (MSO) on PDA medium

Isolate of Pathogenic fungi	Host plant	MSO concentration									
		1.0%		1.5%		2.0%		2.5%		3.0%	
		LG	R	LG	R	LG	R	LG	R	LG	R
<i>Fusarium ox f. sp. lycopersici</i> (Fol2)	Tomato	46.6e	48.2	31.5	65.0	27.0	70.0	0.0	100	0.0	100
<i>F. ox radialis. lycopersici</i> (For15)	Tomato	48.9e	45.6	35.1	61.0	28.1	68.8	0.0	100	0.0	100
<i>Fusarium oxysporum</i> (Fox 1)	Green bean	45.0e	50.0	32.4	64.0	25.0	72.2	0.0	100	0.0	100
<i>F. solani</i> (Fs 5)	Tomato	48.7e	54.8	25.2	72.0	14.2	84.2	0.0	100	0.0	100
<i>F. solani</i> (Fs 3)	Green bean	45.0e	50.0	27.0	70.0	16.2	82.0	0.0	100	0.0	100
<i>Fusarium semibaticum</i> (Fse 2)	Potato	49.6e	44.8	36.0	60.0	28.8	68.0	0.0	100	0.0	100
<i>Alternaria solani</i> (As 3)	Tomato	44.4e	50.6	37.8	58.0	36.6	66.0	0.0	100	0.0	100
<i>Alternaria solani</i> (As 1)	Potato	50.0d	44.0	44.2	50.8	34.0	62.2	0.0	100	0.0	100
<i>Phytophthora infestans</i> (Ph 3)	Tomato	45.0e	50.0	34.0	62.2	27.0	70.0	0.0	100	0.0	100
<i>Phytophthora infestans</i> (Ph 1)	Potato	48.2e	46.4	37.9	57.8	30.6	66.0	0.0	100	0.0	100
<i>Rhizoctonia solani</i> (Rs 5)	Tomato	53.8d	40.2	46.8	48.0	40.3	55.2	25.0	72.2	0.0	100
<i>Rhizoctonia solani</i> (Rs 1)	Green bean	52.2d	42.0	43.2	52.0	36.0	60.0	10.6	88.2	0.0	100
<i>Rhizoctonia solani</i> (Rs 2)	Potato	55.8d	38.0	49.8	44.6	41.4	54.0	27.0	70.0	0.0	100
<i>Sclerotium rolfsii</i> (Sr 5)	Potato	63.0b	30.0	57.6	36.0	52.2	42.0	34.2	62.0	0.0	100
<i>Sclerotium rolfsii</i> (Sr 3)	Green bean	61.0b	32.2	53.6	40.4	46.8	48.0	32.0	64.4	0.0	100
<i>Sclerotina sclerotiorum</i> (Sc 2)	Green bean	58.5c	35.0	53.8	40.2	45.2	49.8	30.6	66.0	0.0	100
<i>Botrytis cinerea</i> (Bc 3)	Green bean	57.7c	35.8	51.4	42.8	44.2	50.8	30.4	66.2	0.0	100
<i>Macrophomina phaseolina</i> (Mph 3)	Green bean	52.2d	42.0	39.6	56.0	34.2	62.0	10.8	88.0	0.0	100
Control		90.0a	0.0	90.0	0.0	90.0	0.0	90.0	0.0	90.0	0.0

LG = colony diameter (mm) . R = Inhibition percentage (%) as compared to the control. Means followed by the same letters are not significantly different according to Duncan's multiple range test ( $P \leq 0.05$ ).

**Table 6.** Reduction % in spores and sclerotia germination of pathogenic fungi on tomato, green bean and potato as affected by different concentration of MSO on PDA medium

Isolate of Pathogenic fungi	Host plant	MSO concentration				
		1.0%	1.5%	2.0%	2.5%	3.0%
Spores germination						
<i>Fusarium ox f. sp. lycopersici</i> (Fol2)	<b>Tomato</b>	51.6	70.0	74.0	100	100
<i>F. ox radicans. lycopersici</i> (For15)	<b>Tomato</b>	50.0	66.2	70.0	100	100
<i>Fusarium oxysporum</i> (Fox 1)	<b>Green bean</b>	55.2	73.4	80.0	100	100
<i>F. solani</i> (Fs 5)	<b>Tomato</b>	57.2	77.8	90.0	100	100
<i>F. solani</i> (Fs 3)	<b>Green bean</b>	54.4	74.4	88.2	100	100
<i>Fusarium semibaticum</i> (Fse 2)	<b>Potato</b>	42.2	60.0	68.2	100	100
<i>Alternaria solani</i> (As 3)	<b>Tomato</b>	<b>54.6</b>	<b>64.4</b>	<b>70.0</b>	<b>100</b>	<b>100</b>
<i>Alternaria solani</i> (As 1)	<b>Potato</b>	52.6	58.0	68.2	100	100
<i>Phytophthora infestance</i> (Ph 3)	<b>Tomato</b>	<b>55.2</b>	<b>64.2</b>	<b>72.2</b>	<b>100</b>	<b>100</b>
<i>Phytophthora infestance</i> (Ph 1)	<b>Potato</b>	53.8	60.2	75.0	100	100
Sclerotia germination						
<i>Rhizoctonia solani</i> (Rs 5)	<b>Tomato</b>	<b>41.0</b>	<b>55.2</b>	<b>71.2</b>	<b>84.0</b>	<b>100</b>
<i>Rhizoctonia solani</i> (Rs 1)	<b>Green bean</b>	<b>44.4</b>	<b>51.4</b>	<b>71.0</b>	<b>80.2</b>	<b>100</b>
<i>Rhizoctonia solani</i> (Rs 2)	<b>Potato</b>	<b>40.4</b>	<b>54.2</b>	<b>70.0</b>	<b>77.4</b>	<b>100</b>
<i>Sclerotium rolfsii</i> (Sr 5)	<b>Potato</b>	35.6	50.8	58.0	64.2	100
<i>Sclerotium rolfsii</i> (Sr 3)	<b>Green bean</b>	32.8	52.4	64.4	70.0	100
<i>Sclerotina. sclerotiorum</i> (Sc 2)	<b>Green bean</b>	42.2	54.8	66.4	74.0	100
<i>Botrytis cinerea</i> (Bc 3)	<b>Green bean</b>	44.6	54.2	70.2	100	100
<i>Macrophomina phaseolina</i> (Mph 3)	<b>Green bean</b>	<b>48.2</b>	<b>57.4</b>	<b>74.0</b>	<b>90.0</b>	<b>100</b>

## Discussion

*Moringa oleifera* Lam. (Family Moringaceae) is commonly known as horseradish tree or drumstick used as phytomedicine such as antioxidant, antimicrobial, anti-inflammatory, antipyretic, antidiabetic, antifertility, antiulcer and antitumor (Fahey, 2005; Foidle *et al.*, 2001; Anwar *et al.*, 2007). In the present study, the antifungal activities of *Moringa oleifera* extracts i.e., MLE, MRE and MSO against 18 plant pathogenic fungi, the causal agents of many root rot, wilt and foliar diseases of tomato, green bean and potato, was investigated in vitro. Our results demonstrated that *Moringa oleifera* leaves extracts (MLE), roots extracts (MRE) as well as moringa seed oil and (MSO) at all tested concentration had antifungal activity against all tested pathogenic fungi. Moringa seed oil (MSO) and MRE showed highly antifungal activities against all tested fungi, as the highest records in linear growth, spore and

sclerotia germination of all tested pathogens were observed with all concentrations of MSO and MRE. The highest recodes of reduction in linear growth, spores and sclerotia germination were observed with all tested isolates of *Fusarium*, *Phytophthora* and *Alternaria*, as they were highly sensitive to MLE, MRE and MSO than all isolates of *Macrophomina*, *Rhizoctonia*, *Botrytis* and *Sclerotium*.

These results are agreement with those obtained by other investigators who found an antifungal activity of moringa plant extracts against several pathogens (Adandonon *et al.*, 2006; Anwar and Rashid, 2007; Al-Asker and Rashed 2010; Abdulmoneim *et al.*, 2011; Moyo *et al.*, 2012; Talreia, 2010; Seint and Masara, 2011, El-Mohamedy and Aboelfetoh, 2014 a,b). Jed and Fahey (2005) noted that antimicrobial activity of *Moringa oleifera* extracts may be attributed two main phytochemicals viz., pterygospermin and benzyl isocyanate which have strong antifungal and antimicrobial activity. In this respect, many essential oils and plant extracts have been found to be potent fungitoxic agents against many plant pathogens (Satish *et al.*, 2007; Jamil *et al.*, 2007; Anwar and Rashid, 2007; Sun *et al.*, 2007, Siripornvisal and Ngamchawee, 2010; El-Mohamedy *et al.*, 2013; Tabassum and Vidyasagar, 2013; Abd el-kader *et al.*, 2013).

However, the harmful effects on fungi were restricted in: (a) partial or complete inhibition on spore germination, sporulation or mycelia growth and (b) alternation in physiology and biochemistry activities of the fungal cells. In additions, secondary compounds, considered as final products of plant metabolism or metabolite refuses, have important ecological functions for the plant which synthesize them. One of these functions is to protect the plants against infection by pathogens (Price, 2000; Anwar and Rashed, 2007). Therefore, much plant extracts exhibited inhibitory properties in challenge tests against microorganisms. These extracts, however, contained specific component that can inhibit the growth of certain microorganisms (Bowers and Locke, 2000; Dubey *et al.*, 2009; Jamil *et al.*, 2010; Moyo *et al.*, 2013).

The fungicidal effect of Moringa extracts on some soil-borne fungi such as *Rhizoctonia*, *Pythium* and *Fusarium* was recorded by many investigators (Chuang *et al.*, 2007; Al-Asker and Rashed 2010; Raj *et al.*, 2011; Moyo *et al.*, 2012; Hadi and Klefi, 2013; El-Mohamedy and Aboelfetoh, 2014 a, b). Dwivedi and Enespa (2012) indicate that *Moringa oleifera* extracts (leaves, bark and seeds) 75 % (v/v) showed significant inhibition in the mycelial growth of *Fusarium solani* and *Fusarium oxysporum* f. sp. *lycopersici*. Raj *et at.* (2011) demonstrated that aqueous extract of *Moringa oleifera* (Lam.) Root showed maximum inhibition against many pathogenic fungi and bacteria in vitro. They also noted that the phytochemical screening revealed the presence of alkaloids,

flavonoids, saponins, erpenoids, steroids, tannins, cardioglycosides, aminoacids and proteins which have antifungal and antibacterial activities.

Studies on spore germination represent an integral part of the ecological studies of the pathogenic fungi, as spores are the specialized structures capable of initiating new growth. Our studies also demonstrated that spore/sclerotia germination of all tested pathogens were gradually decreased to reach 100% reduction with increasing concentrations of MLE (50%), MRE(20%) and MSO (3%). Meanwhile, sclerotia germination of *R. solani*, *S. rolfsii* and *M. phaseolina*, *S. sclerotiorum* and *B.cinerea* showed less sensitive against MLE, MRE and MSO in ascending order. These findings are in agreement with those reported by many investigators (Bowers and Locke, 2000; Dwivedi and Enespa, 2012; Abdulmoneim *et al.*, 2011; Chuang *et al.*, 2007, El-Mohamedy and Aboelfetoh, 2014 a, b)

## Conclusion

The results of this study demonstrated that, Moringa leaves extract MLE, Moringa roots extract MRE and Moringa seed oil MSO were the most efficient and might be as promising natural compounds for controlling such plant pathogenic fungi. Therefore, it could be used as natural fungicide to compact pathogenic fungi and thus reduced the dependence on the synthetic fungicides. More studies are still needed in the future to test the antifungal activities of moringa extracts on other different fungal plant diseases *in vitro* and under field conditions.

## Acknowledgments

This work was financially supported by The Affairs of Research Projects, National Research Centre, Egypt. Grand NO. 110-30-138 (2016-2019).

## References

- Abdulmoneim, M., Saadabi, M. and Abu Zaid, I. E. (2011). An *In vitro* antimicrobial activity of *Moringa oleifera* L. seed extracts against different groups of microorganisms. Australian Journal of Basic and Applied Sciences 5:129-134.
- Agrios, G. M. (2005). Plant Pathology. 5th ed. New York: Academic Press.
- Al-Askar, A. A. and Rashad, Y. M. (2010). Efficacy of some plant extracts against *Rhizoctonia solani* on pea. Journal of Plant Protection Research 50:239-243.
- Anwar, F. and Rashid, U. (2007). Physico-chemical characteristics of *Moringa Oleifera* seeds and seed oil from a wild provenance of Pakistan. Pakistan Journal of Botany 39:1443-1453.

- Anwar, F., Latif S., Ashraf, M. and Gilan, A. H. (2007). *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytotherapy Research* 21:17-25.
- Ashfaq, M., Basra, S. M. A. and Ashfaq, U. (2012). Moringa: A miracle plant of agro-forestry. *Journal of agriculture and social sciences* 8:115-122.
- Bowers, J. H. and Locke, J. C. (2000). Effect of botanical extracts on the population density of *Fusarium oxysporum* in soil and control of *Fusarium* wilt in the greenhouse. *Plant Disease* 84:300-305.
- Chuang, P. H., Lee, C. W. Chou, J. Y, Murugan, M., Shieh, B. J. and Chen, H. M. (2007). Antifungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. *Bioresource Technology* 98:232-236.
- Co Stat (2005). Cohort Software, 798 Lighthouse Ave. PMB 320 Monterey, USA, pp. 65.
- Duncan, D. B. (1955). Multiple range and multiple F test. *Biometrics* 11:1-24.
- Dubey, R. C, Kumar, H. and Pandey, R. R. (2009). Fungitoxic effect of neem extracts on growth and sclerotial survival of *Macrophomina phaseolina in vitro*. *Journal of American Science* 5:17-24 .
- Dwivedi, S. K. and Enespa, A. (2012). Effectiveness of extract of some medical plants against soil borne fusaria causing diseases on *Lycopersicon esculantum* and *Solanum melongena*. *Int. International Journal of Pharma and Bio Sciences* 3:1171-1180.
- El-Mohamedy, R. S. R., Shafeek, M. R., Abd El-Samad, E. H., Salama, D. M. and Rizk, F. A. (2017). Field application of plant resistance inducers (PRIs) to control important root rot diseases and improvement growth and yield of green bean (*Phaseolus vulgaris* L.) *Australian journal of crop science* 11:496-505.
- El-Mohamedy, R. S. R., Aboelfetoh, M. A. and Safia, M. A. (2014). Preliminary studies on response of *Moringa oleifera* plants to infection with some soil borne plant pathogenic fungi. *International Journal of Current Microbiology and Applied Sciences* 3:389-397.
- El-Mohamedy, R. S. R. and Aboelfetoh, M. A. (2014a). Evaluation of antifungal activity of *Moringa oleifera* extracts as natural fungicide against some plant pathogenic fungi *In-vitro*. *Journal of Agricultural Technology* 10:963-982.
- El-Mohamedy, R. S. R. and Aboelfetoh, M. A. (2014b). Antifungal activity of *Moringa oleifera* oil and seed extract against some plant pathogenic fungi. *Middle East Journal of Agriculture Research* 3:242-249.
- El-Mohamedy, R. S. R., Abdel-Kader, M. M., Abd-El-Kareem, F. and El-Mougy, N. S. (2013). Essential oils, inorganic acids and potassium salts as control measures against the growth of tomato root rot pathogens *in vitro*. *Journal of Agricultural Technology* 9:1507-1520.
- Fahey, Sc. D. (2005). *Moringa oleifera* A review of the medical evidence for its nutritional, therapeutic and prophylactic properties, Part1. <http://www.TFLJournal.org>. 1:1-15.
- Fawzi, E. M., Khalil, A. A. and Afifi, A. F. (2009). Antifungal effect of some plant extracts on *Alternaria alternata* and *Fusarium oxysporum*. *African Journal of Biotechnology* 8:2590-2597.
- Foidle, N., Makkar, H. P. S. and Becker, K. (2001). The potential of *Moringa oleifera* for agricultural and industrial uses. *In: Fugile, L.J., (ed.), The Miracle Tree: The Multiple Attribute of Moringa*. pp. 45-76.
- Garima, M., Pradeep, S., Ramesh, V., Sunil, K., Saurabh, S., Jha, K. K. and Khosa, R. L. (2011). Traditional uses, phytochemistry and pharmacological properties of *Moringa oleifera* plant: An overview. *Der Pharmacia Letter* 3:141-164.
- Gnanamanickam, S. S. (2002). *Biological Control of Crop Diseases*. New York. Basel: Marcel Dekker, Inc. pp. 15.

- Lee, S. O., Gyung, J. C., Kyoung, S. J., He Kyoung, L., Kwang, Y. C. and Jin-Cheol, K. (2007). Antifungal Activity of five plant essential oils as fumigant against postharvest and soil borne plant pathogenic fungi. *The Plant Pathology Journal* 23:97-102
- Hadi, M. and Kashefi, B. (2013). Study on effect of some medicinal plant extracts on growth and spore germination of *Fusarium oxysporum* schlecht. *In vitro American-Eurasian Journal of Agriculture and Environmental Sciences* 13:581-588.
- Jamil, A., Shahid, M., Khan, M. M. and Ashraf, M. (2010). Screening of some medicinal plants for isolation of antifungal proteins and peptides. *Pakistan Journal of Botany* 39:211-221.
- Jed, W. and Fahey, S. D. (2005). *Moringa oleifera* : A review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. Part 1 *Trees for Life Journal*. pp. 1-5.
- Moyo, B., Masika, P. J. and Muchenje, V. (2012). Antimicrobial activities of *Moringa oleifera* Lam leaf extracts, *African Journal of Biotechnology* 11:2797-2802.
- Price, M. L. (2000). The *Moringa* Tree purification and activity assay of the coagulant protein from *Moringa oleifera* seed. *Water Research* 39:2338-2344.
- Rai, M. and Carpinella, M. (2006). *Naturally Occurring Bioactive Compounds*. Elsevier, Amsterdam. p. 502.
- Price, M. L., (2000). The *Moringa* Tree purification and activity assay of the coagulant protein from *Moringa oleifera* seed. *Water Research* 39:2338-2344.
- Raj, A. J., Gopalakrishnan, V. K., Yadav, S. A. and Dorairaj, S. (2011). Antimicrobial Activity of *Moringa oleifera* (Lam.) Root Extract. *Journal of Pharmacy Research* 4:1426 -1430.
- Satish, S., Mohana, D. C., Raghavendra, M. P. and Raveesha, K. A. (2007). Antifungal activity of some plant extracts against important seed borne pathogens of *Aspergillus* sp. *Journal of Agricultural Technology* 3:109-119.
- Seema, M. S. S., Sreenivas, N. D. and Devaki, N. S. (2011). *In vitro* studies of some plant extracts against *Rhizoctonia solani* Kuhn infecting FCV tobacco in Karnataka Light Soil, Karnataka, India. *Journal of Agricultural Technology* 7:1321-1329.
- Siripornvisal, S. and Ngamchawee, K. (2010). Utilization of herbal essential oils as biofumigant against fungal decay of tomato during storage. *Proceedings 16th Asian Agricultural Symposium, Bangkok, Thailand*. pp. 655-658.
- Sun, O. L., Gyung, J. C., Kyoung, S. J., He, K. L., Kwang, Y. C. and Jin-Cheol, K. (2007). Antifungal Activity of Five Plant Essential Oils as Fumigant against Postharvest and Soil borne Plant Pathogenic Fungi. *The Plant Pathology Journal* 23:97-102.
- Tabassum, N. and Vidyasagar G. M. (2013). Antifungal investigations on plant essential oils, A review. *International Journal of Pharmacy and Pharmaceutical Sciences* 5:19-28.
- Winer, B. J. (1971). *Statistical Principles in Experimental Design*. In 2nd ed. New York: MiGraw-Hil.
- Zaker, M. (2016). Natural Plant Products as Eco-friendly Fungicides for Plant Diseases Control- A Review. *The Agriculturists* 14:134-141.

(Received: 11 May 2018, accepted: 30 June 2018)