
Nano-particles from *Chaetomium brasiliense* to control *Phytophthora palmivora* caused root rot disease in durian var Montong

Tongon, R.^{*1}, Soyotong, K.¹, Kanokmedhakul, S.² and Kanokmedhakul, K.²

¹Department of Plant Production Technology, Faculty of Agricultural Technology, King Mongkut's Institute of Technology Ladkrabang, Bangkok, Thailand; ²Department of Chemistry and Center for Innovation in Chemistry, Faculty of Science, Khon Kaen University, Khon Kaen, Thailand.

Tongon, R., Soyotong, K., Kanokmedhakul, S. and Kanokmedhakul, K. (2018). Nano-particles from *Chaetomium brasiliense* to control *Phytophthora palmivora* caused root rot disease in durian var Montong. International Journal of Agricultural Technology 14(7): 2163-2170.

Abstract The antagonistic fungus, *Chaetomium brasiliense* was tested to control *Phytophthora palmivora* causing rot disease of Durian (*Durio zibithenus* L.) var Montong by crude extracts and Nano-particles derived from *Ch. brasiliense*. Crude extracts of antagonistic fungus was tested for antifungal biological activities. The crude extracts from antagonistic fungus with hexane, ethyl acetate and methanol were tested against *P. palmivora*. Crude ethyl acetate from *Ch. brasiliense* gave significantly against *P. palmivora* which the ED₅₀ values was 17.46 µg/ml. Testing Nano-particles were tested for antifungal activities. The results showed nano - particles from *Ch. brasiliense* gave effectively significantly inhibition of colony growth and spore production which the ED₅₀ values were 1.08 µg/ml, and 8.68 µg/ml respectively. Application of Nano - particles to control the *P. palmivora* causing root rot disease of durian in pot experiment was successfully done. The results showed nano-particles from *Ch. brasiliense* reduced the root rot disease on durian of 40%. The nano-particles from *Ch. brasiliense* gave significantly high plant growth which were 79.5 cm when compared to the non-treated control.

Keywords: Nano-particles, *Chaetomium brasiliense*, *Phytophthora palmivora*, biological control

Introduction

Durian (*Durio zibithenus* L) is originated from the region of Borneo and Sumatra, growing wild in the Malay peninsula, cultivated in a wide region from India to New Guinea four hundred years ago. It was across to Myanmar, and cultivated through Thailand and South Vietnam (Morton, 2000). The problem for durian cultivation in Thailand faced the root rot which caused by *Phytophthora* spp. It can infect all stages of growing durian trees. The symptoms are appeared as root rot, leaves and stem blight, bark and fruit rot. Chemical fungicides have

* **Coressponding Author:** Tongon, R.; **Email:** rutt1409@gmail.com

been used to control this disease and tended to lead the negative side effect to the environment. In addition, *Phytophthora* spp become resistant to those fungicide application eg. metalaxyl and related compounds (Erwin and Ribeiro, 1996). Tye eco-friendly management of *Phytophthora* diseases would be done to reduce the application cost. The application of biocontrol agents against *Phytophthora* rot has become an importance research aspect and is being carried out all over the world (Naqvi, 2004).

Biological control of plant pathogen is known as using antagonists against plant pathogens. The antagonism is the control mechanism to reduce the pathogen onocullum and disease incidence. Antagonism is expressed between two organisms including antibiosis, competition and parasitism (Cook and Baker, 1983). However, there are many reports indicated that the antagonists can express to against several plant pathogens eg such as *Chaetomium* spp; and *Emericella nidulans*. (Xu *et al.*, 2017; Hung, *et al.*, 2015a; Arjona-Girona *et al.*, 2018; Tran *et al.*, 2007; Song *et al.*, 2017).

The objective was to test the nano-particles derived from *Ch. brasiliense* to inhibit *Phytophthora* sp causing root rot in durian var Montong.

Materials and methods

Isolation of pathogen and testing pathogenicity

Phytophthora sp was isolated the durian diseased plant parts eg bark and root rot using tissue transplanting method. The collected diseased samples were washed with sterilized water, and cut it into small pieces, soaked 1% clorox for 3 min. The piece samples of diseased part was moved to water agar (WA), then incubated at room temperature to observe growing colonies, then gently moved to PDA to get pure culture. Morphology was identified and observed the characters of fungus under compound microscope. The pathogenicity was tested using detached leaves technique. The agar plugs of *Phytophthora* sp were transferred to wounded leaves. The non-inoculated leaves with sterilized agar discs was done to serve as controls, and incubated at room temperature (27-30 °C) for seven days. The experiment was replicated four times. Disease incidence was calculated as number of infected plants/ total number of tested plants x 100, and disease ratings was evaluated as 0= healthy plants, and 3= seriously infected plants (Soytong, 2010).

Morphological study of Chaetomium brasiliense

Chaetomium brasiliense used in this research study is offered from Assoc. Prof. Dr. Kasem Soytong, Department of Plant Production Technology, Faculty

of Agricultural Technology, King Monkut's Institute of Technology Ladkrabang (KMITL), Bangkok, Thailand. The culture was grown on potato dextrose agar and observed morphological characters under compound microscope.

Dual culture test

The experimental design was used as Completely Randomization with four replications. Dual culture was done by followed the methods of Soyong (1992). *Ch. brasiliense* and *Phytophthora* sp. were separately cultured on PDA at room temperature (27-30 °C) for seven days. The 0.5 cm diameter of sterilized cork borer was cut at the peripheral colony in each culture and moved to PDA plates (9 cm). The agar plug of *Phytophthora* sp. was moved to PDA plate on one side of the plate that opposited the agar plug of *Ch. brasiliense*. The plug of each fungus was transferred to PDA plate served as the controls. All plates were incubated at room temperature for 30 days. Colony diameter (cm) and conidia production in the dual culture plates and control plates were recorded. The number of conidia was counted by using haemocytometer.

The inhibition of colony growth or conidia production were calculated using the following formula:

$$\% \text{ inhibition} = (A-B) / A \times 100$$

Where, A was the colony diameter or number of conidia produced by the pathogen on the control plates and B was the colony diameter or number of conidia produced by the pathogen in the dual culture plate.

Testing crude extracts of *Chaetomium brasiliense* to against *Phytophthora* spp in vitro

Crude extracts of *Ch. brasiliense* were evaluated to inhibit *Phytophthora* sp. The experimental design was conducted using factorial experiment in Completely Randomization, and repeated four times. Factor A was crude hexane, crude ethyl acetate and crude methanol. Factor B was 0, 10, 50, 100, 500, and 1,000 µg/ml. Each crude extract was poured to 2% dimethyl sulphite and mixed to 30 ml potato dextrose agar, then autoclaved at 121°C, 15 lbs/inch² for 35 minutes. *Phytophthora* sp colony was cut at peripheral colony with sterilized cork borer (0.5 mm). Agar plug of pathogen was transferred to the middle of PDA media in plate incorporated with each and incubated at room temperature (27-30°C) until the pathogen growing full plate. Data were recorded as colony diameter, number of conidia. The pathogen colony growth and conidia production were calculated the inhibition using the above formula. Data were statistically computed analysis of variance. Means were compared using Duncan Multiple's Range Test. The effective dose was computed using probit analysis.

In vitro testing nano-particles from Chaetomium brasiliense to control root rot disease in durian var Montong

Chaetomium brasiliense was cultured in potato dextrose broth (PDB) at room temperature (27-30 C) for 30 days. The biomass was filtered through cheesecloth and air-dried. Fresh and dried biomass were weighted. The biomass was ground in electrical blender, then extracted by adding the same volume of hexane, and kept in stationary phase for 7 days at room temperature. The filtrate was get it through Whatman filter paper, evaporated in rotary vacuum evaporator to yield crude hexane. The remaining marc was consecutively extracted with ethyl acetate and methanol to get crude extracts using the same procedure as hexane to yield crude ethyl acetate (EtOAc) and crude methanol (MeOH). The nano- particles were done by followed the method of Dar and Soyong (2014), to yield Nano-CBH, Nano-CBE and Nano-CBM. Those nano-particles were evaluated to inhibit *Phytophthora* sp. The experimental design was used two factors factorial experiment in Completely Randomization. The experiment was repeated four times. Factor A was Nano-CBH, Nano-CBE and Nano- CBM. Factor B was 0, 1, 5 and 10 µg/ml. Each Nano-particle was dissolved in 2% dimethyl sulfoxide, and mixed to PDA before autoclaved at 121°C, 15 lbs/inch² for 30 min. The culture of *Phytophthora* sp was cut at peripheral colony with sterilized cork borer (0.5 mm). Agar plug of *Phytophthora* sp was moved to the middle of PDA mixed with each nano-particles. All plates were incubated at room temperature until the pathogen in control growing full plate. The pathogen from each treatment were observed abnormal spores under compound microscope. Data were stisticaally computed. The effective dose was calculated using probit analysis.

Results

Pathogenicity test

Pathogenicity test was conducted by detached leaves method which resulted durian leave var Montong showed brown hydrolysis expand around agar plug of pathogen. In control, leaves remained healthy as seen in Fig.1.

Dual culture test

Ch. brasiliense was proved it ability to inhibit plant pathogen *P. palmivora* causing disease of durian by using dual culture tests. The results showed that *Ch. brasiliense* gave significantly growth inhibition of *P. palmivora* which were 58.33% and showed significantly inhibited the spore production of pathogen of 88.82% (Table 1).



Figure1. Pathogenicity test of *Phytophthora* sp on Durian leaves

Testing crude extracts of Chaetomium brasiliense to against Phytophthora spp in vitro

The crude extracts from *Ch. brasiliense* were used to test their abilities to control the growth of *P. palmivora*. The results showed that crude hexane extract gave inhibition of *P. palmivora* colony growth which was 57.75% at the concentration of 1,000 $\mu\text{g/ml}$ with the ED_{50} values of 204.28 $\mu\text{g/ml}$ when compared to the control. Crude ethyl acetate (EtOAc) extract and showed significantly highest inhibition for the spore production of *P. palmivora* as 97.72% at the concentration of 1,000 $\mu\text{g/ml}$. which the ED_{50} values were 182.05 $\mu\text{g/ml}$, followed by crude hexane extract and methanol which the ED_{50} values were 538.95 and 722.08 $\mu\text{g/ml}$., respectively (Table 2).

Table 1. Colony growth and number of spore on antagonistic dual-culture tests

Antaginistic fungi	<i>P. palmivora</i>			
	Colony diameter (cm)	Growth inhibition (%)	Spore number ($10^6/\text{ml}$)	Spore inhibition (%)
Control	9.00 ^{a1}	-	27.43 ^a	-
<i>Ch. brasiliense</i>	3.75 ^b	58.33 ^b	5.31 ^b	88.82 ^a
C.V. (%)	7.90		19.39	

¹ Means of four repeated experiments. Means followed by the same letters are not significantly differed by DMRT at $P=0.01$.

In vitro testing nano-particles from Chaetomium brasiliense to control Phytoththora sp causing durian rot var Montong

Result from nano-particles showed high efficacy antifungal activity of nanoparticles from *Ch. brasiliense* against *P. palmivora*. The result of Nano particles from *Ch. brasiliense* was showed that Nano-CBH, Nano-CBE and Nano-CBM gave highly significant inhibited the colony growth of *P. palmivora* as 90.00% which the ED_{50} values of 1.25, 1.12, 1.08 $\mu\text{g/ml}$., respectively and gave highly significant inhibition for the spore production of *P. palmivora* as 100% which the ED_{50} values of 8.68, 12.20 and 10.77ppm (Table 3).

Table 2. Crude extracts of *Ch. brasiliense* testing for growth inhibition of *P. palmivora*

Crude metabolites	Concentration (ppm)	Colony diameter (cm) ^{2,3}	Growth inhibition (%) ^{2,3}	ED ₅₀ (µg/ml)	Number of spores ^{2,3} (10 ⁵)	Spore Inhibition (%) ^{2,3}	ED ₅₀ (ppm)
CB-H	0	5.00 ^a	-		32.25 ^a	-	
	10	5.00 ^a	0.00 ^g		29.00 ^b	10.47 ^{fg}	
	50	5.00 ^a	0.00 ^g	770.60	27.25 ^b	15.34 ^{fg}	538.95
	100	5.00 ^a	0.00 ^g		21.25 ^d	33.77 ^e	
	500	3.29 ^c	34.00 ^e		13.75 ^e	57.29 ^c	
	1000	2.02 ^e	57.75 ^c		11.00 ^f	65.69 ^b	
CB-E	0	5.00 ^a	-		32.25 ^a	-	
	10	5.00 ^a	0.00 ^g		28.50 ^b	11.35 ^{fg}	
	50	5.00 ^a	20.00 ^f	204.28	26.50 ^b	17.69 ^f	182.05
	100	3.37 ^e	32.5 ^e		15.25 ^e	47.44 ^d	
	500	1.25 ^f	75.00 ^b		10.00 ^f	96.25 ^a	
	1000	0.86 ^g	82.75 ^a		0.50 ^g	97.72 ^a	
CB-M	0	5.00 ^a	-		32.25 ^a	-	
	10	5.00 ^a	0.00 ^g		29.25 ^{ab}	9.04 ^g	
	50	5.00 ^a	0.00 ^g		27.75 ^b	13.77 ^{fg}	722.08
	100	5.00 ^a	0.00 ^g	-	21.00 ^d	34.33 ^e	
	500	4.12 ^b	17.5 ^f		21.50 ^d	33.11 ^e	
	1000	2.65 ^d	47.00 ^d		11.25 ^f	65.23 ^b	
C.V. (%)		3.96	9.52		7.13	14.92	

¹ Means of four repeated experiments. Means followed by the same letters are not significantly differed by DMRT at P=0.01.

Table 3. Nano particle extracts of *Ch. brasiliense* testing for growth inhibition of *P. palmivora*

Nano particles	Concentration (ppm)	Colony diameter (cm) ^{2,3}	Growth inhibition (%) ^{2,3}	ED ₅₀ (µg/ml)	Number of spores ^{2,3} (10 ⁵)	Spore Inhibition (%) ^{2,3}	ED ₅₀ (ppm)
Nano CBH	0	5.00 ^a	-		32.37 ^a	-	
	1	2.75 ^b	45.00 ^e		13.25 ^b	59.20 ^f	
	3	1.41 ^e	71.75 ^{cd}	1.25	3.50 ^d	89.05 ^c	8.68
	5	1.35 ^{ef}	73.00 ^{cd}		1.50 ^e	95.42 ^b	
	10	1.11 ^g	77.75 ^{bc}		0.50 ^e	98.50 ^{ab}	
	15	0.5 ^h	90.00 ^a		0.00 ^e	100.00 ^a	
Nano CBE	0	5.00 ^a	-		32.37 ^a	-	
	1	2.60 ^{bc}	47.50 ^e		10.00 ^c	69.12 ^e	
	3	1.67 ^d	66.50 ^d	1.12	1.50 ^e	95.40 ^b	12.20
	5	1.17 ^{fg}	76.50 ^{bc}		0.25 ^e	99.21 ^a	
	10	1.00 ^g	80.00 ^b		0.00 ^e	100.00 ^a	
	15	0.5 ^h	90.00 ^a		0.00 ^e	100.00 ^a	
Nano CBM	0	5.00 ^a	-		32.37 ^a	-	
	1	2.50 ^c	50.00 ^e		8.75 ^c	72.94 ^d	
	3	1.52 ^{de}	69.50 ^d		0.75 ^e	97.79 ^{ab}	
	5	1.12 ^{fg}	77.50 ^{bc}	1.08	0.25 ^e	99.21 ^a	10.77
	10	0.5 ^h	90.00 ^a		0.00 ^e	100.00 ^a	
	15	0.5 ^h	90.00 ^a		0.00 ^e	100.00 ^a	
C.V. (%)		8.74	7.90		13.90	2.71	

¹ Means of four repeated experiments. Means followed by the same letters are not significantly differed by DMRT at P=0.01.

Discussion

Morphological study was done in previous experiment to prove identification of *P. palmivora* which cultured on potato dextrose agar. The colony is white, slow growing, non septate mycelia, sporangia produce readily and abundantly. The identity in morphological characteristics are consistent with descriptions of *Phytophthora palmivora* (Erwin and Ribiero, 1996).

The results showed that crude ethyl acetate (EtOAc) extract from *Ch. brasiliense* gave highest inhibition of *P. palmivora* colony growth which was 57.75% at the concentration of 1,000 µg/ml with the ED₅₀ values of 204.28 ppm when compared to the control. As similar in the previous report, found that crude extract of *Ch. globosum* CG05, *Ch. cupreum* CC3003, *Ch. lucknowense* CL01 showed ability to inhibit mycelial growth and spore production of *P. palmivora* PHY02 in laboratoty test (Hung *et al.*, 2015b). This result is also similar reported by Hung *et al.*, (2015a) found that crude extracts of these *Chaetomium* species exhibited antifungal activities against mycelial growth of *P. nicotianae*, with effective doses of 2.6~101.4 µg/ml.

It was found that nano-particles derived from *Ch. brasiliense* was actively inhibited the *P. palmivora*. Nano-CBH, Nano-CBE and Nano-CBM gave highly significant inhibition of the colony growth of *P. palmivora* at 90.00% and gave highly significant inhibition for the spore production of *P. palmivora* at 100%. This result is similar reported by Thongkham (*et al.*, 2017) found that nano-particles derived from *Ch. cupreum* to inhibit mycelial growth and spore production of *Phytophthora* spp causing root rot in durian. This study was also similar to Dar *et al.* (2013) that nano particles of *Ch. globosum* and *Ch. cuperum* were proved to againt *F. oxysporum* f.sp. *lycopersici* and *Colletotrichum capsici*.

Acknowledgement

I would like to acknowledge the financial support from Thailand Research Fund (Grant No RTA5980002).

References

- Arjona-Girona, I. and López-Herrera, C. J., (2018). Study of a new biocontrol fungal agent for avocado white root rot. *Biological Control*. 117:6-12.
- Cook, R. J. and Baker, K. F. (1983). The nature and practice of biological control of plant pathogen. American Phyto pathological Society, St. Paul, MN. 539.
- Dar, J. and Soyong, K. (2014) Construction and characterization of copolymer nanomaterials loaded with bioactive compounds from *Chaetomium* species. *Journal of Agricultural Technology*. 10:823-831.

- Dar, J. and Soyong, K. (2013). In-vitro testing of nanomaterials containing globosum ethyl acetate extract against *Fusariumoxysporum* f. sp. *lycopersici* (race 2). International Conference on Integration of Science and Technology for Sustainable Development 2013, KMITL, Bangkok, Thailand November 28-29, 2013. pp. 5.
- Erwin, D. C. and Ribeiro, O. K. (1996). Phytophthora diseases worldwide. St. Paul: APS Press.
- Hung, P. M., Wattanachai, P., Kasem, S. and Poaim, S. (2015a). Efficacy of *Chaetomium* species as biological control agents against *Phytophthora nicotianae* root rot in citrus. *Mycobiology*. 43:288-296.
- Hung, P.M., Wattanachai, P., Kasem, S. and Poaim, S. (2015b). Biological control of *Phytophthora palmivora* causing root rot of pomelo using *Chaetomium* spp. *Mycobiology*. 43:63-70.
- Morton, J. F. (200). Fruits of warm climates. Echo Point Books and Media. Brattleboro, United States. pp. 505.
- Naqvi, S. A. (2004). Diagnosis and management of certain important fungal diseases of citrus. In: Naqvi SA, editor. Diseases of fruits and vegetables. Dordrecht: Kluwer Academic Publishers. pp. 247-290.
- Song, J., Soyong, K. and Kanokmedhakul, S., (2017). Antifungal activity of *Emericella nidulans* against *Pyricularia oryzae* causing rice blast. *Journal of Agricultural Technology*. 13:1205-1209.
- Soyong, K. (2010). Evaluation of *Chaetomium*-biological fungicide to control *Phytophthora* stem and root rot of durian. *Research Journal*. 3:117-124.
- Thongkham, D., Soyong, K. and Kanokmedhakul, S. (2017). Efficacy of nano particles from *Chaetomium cupreum* to control *Phytophthora* spp. causing root rot of durian. *Journal of Agricultural Technology*. 13:1295-1300.
- Tran, H., Ficke, A., Asiimwe, T., Höfte, M. and Raaijmakers, J. M. (2007). Role of the cyclic lipopeptide massetolide A in biological control of *Phytophthora infestans* and in colonization of tomato plants by *Pseudomonas fluorescens*. *New Phytologist*. 175:731-742.
- Xu, T., Li, Y., Zeng, X., Yang, X., Yang, Y., Yuan, S., Hu, X., Zeng, J., Wang, Z., Liu, Q. and Liu, Y., (2017). Isolation and evaluation of endophytic *Streptomyces endus* OsiSh-2 with potential application for biocontrol of rice blast disease. *Journal of the Science of Food and Agriculture*. 97:1149-1157.

(Received: 28 August 2018, accepted: 27 October 2018)