Effects of four year sewage sludge application on some morphological traits and chlorophyll content in basil (*Ocimum basilicum* L.)

Azadeh kashani^{1*}, Hemmatollah Pirdashti² and Keyan Kashani³

¹Faculty of Crop Sciences, Sari Agricultural Sciences and Natural Resources University, Sari, Iran, ²Genetic and Agricultural Biotechnology Institute of Tabarestan, Sari Agricultural Sciences and Natural Resources University, Sari, Iran, ³Department of Biology, Islamic Azad University, Sari Branch, Iran

Azadeh Kashani, Hemmatollah Pirdashti and Keyan Kashani (2013) Effects of four year sewage sludge application on some morphological traits and chlorophyll content in basil (*Ocimum basilicum* L.). International Journal of Agricultural Technology 9(2): 451-460.

Abstract In order to study the effect of four year enriched and none-enriched sewage sludge application on some morphological traits and chlorophyll contents of basil, an experiment was conducted in randomized complete blocks design with three replications in 2009. Treatments included control (no fertilizers used), chemical fertilizers (urea, potassium sulfate and super phosphate triple 70, 60 and 50 kg ha⁻¹, respectively) and sewage sludge (20 and 40 Mg ha⁻¹ alone and plus 50% chemical fertilizer). Results showed that different amounts of fertilizer treatments had a significant effect on the studied traits. The maximum inflorescence number, inflorescence length and dry matter yield obtained by using none-enriched sewage sludge as the rate of 20 Mg ha⁻¹. Maximum plant height, stem diameter, chlorophyll b and chlorophyll a+b obtained by application of 20 Mg ha⁻¹ sewage sludge markedly decreased, most of the studied traits where as the minimum leaf area index (LAI) was obtained in both control and 40 Mg ha⁻¹ sewage sludge plus 50% chemical fertilizer. Thus, applying 20 Mg ha⁻¹ sewage sludge either alone or plus 50% chemical fertilizer, had the best effect on studied traits compared to both control and CF treatments.

Key words: Basil, Chemical fertilizer, Chlorophyll, Dry matter yield, Sewage sludge.

Introduction

Sewage sludge is a rich source of elements that are needed for plants and therefor it is considered as a cheap fertilizer by farmers (Singh and Agraval, 2008). Solid phase of sewage sludge consists of large amounts of elements that are needed by plants like nitrogen, phosphorus and elements and so using them on farms is economically important. According to Magdoff and Weil (2004)

^{*} Corresponding author: Azadeh kashani; e-mail: kashani.azadeh@gmail.com; pirdasht@yahoo.com

and Shober *et al.* (2003) the first important element in application of sewage sludge is nitrogen that's been reported between 0 to 56%, but sewage sludge can supply many essential macro and micro elements for plants.

Spnmez and Bozkurt (2006) found that application of sewage sludge can increase the amount of nitrogen, phosphorus, potassium and magnesium in lettuce leaves (*Lactuca sativa*). Warman and Termeer (2005) also reported that sewage sludge contains elements like iron, magnesium, cooper, molybdenum and chlorine which essentially needed by most plants. However, sewage sludge contains a great amount of salt that increases the electrical conductivity of soil. High presence of elements such as sodium and calcium is the reason for increase in sewage sludge affects the crops yield differently. Haghighi (2011) showed that adding sewage sludge to soil causes a significant increase in dry matter yield in celery, rather than spinage and lettuce. Today medicinal plants cultivation is an important branch of agriculture that is used to extract and production of basic material for many drugs. Usage of medicinal plants in developed countries is highly increasing, as 90% of people in these countries use herbal medicines (Akbarnejad *et al.*, 2010).

Basil (*Ocimum basilicum* L.) as a medicinal plant, is of particular importance and it is used for treatment of some heart diseases and migraine and nervous headaches. Since global approaches in production of medicinal plants are going towards sustainable agriculture solutions and management practices like application of organic fertilizers in order to improve quantitative and qualitative yield (Ferreras *et al.*, 2006; *Darzi et al.*, 2008). Recently in Iran some researches has been done on chemical aspects and pollution of sewage sludge on physiological characteristics of medicinal plants. Therefore, this research has been performed to study the long-term application effect of sewage sludge and chemical fertilizer on chlorophyll and physiological parameters in basil (*Ocimum basilicum* L.)

Materials and methods

This experiment has been conducted in research farms of Sari Agricultural Science and Natural Resources University in 2009, Iran, located in north of Iran (34°33'N; 52°6'E and altitude 16 m). The characteristics of soil and non-enriched sewage sludge used in this study are presented in Table 1. The soil of the experiment site was silty clay (10.33% sand, 46.33% clay and 43.33% silt) with 2.43% organic matter. Experiment was arranged in randomized complete block design with six treatments and three replications. Treatments included control (no fertilizers used), chemical fertilizer (urea,

potassium sulphate, super phosphate triple, 70, 60 and 50 kg ha⁻¹ respectively), SS_{20} : sewage sludge 20 Mg ha⁻¹, $SS_{20}+\frac{1}{2}$ CF: sewage sludge 20 Mg ha⁻¹+50% chemical fertilizer (NPK:35-30-25), SS₄₀: sewage sludge 40 Mg ha⁻¹, SS₄₀+¹/₂ CF: sewage sludge 40 Mg ha⁻¹+50% chemical fertilizer (NPK:35-30-25). All treatments has been added to experimental plots during 2006-2009 (four continuous years). The sewage sludge and fertilizer P and K were applied as pretreatment before planting in the plots $(3 \times 1/5 \text{ m}^2)$ and then incorporated in soil. Nitrogen was applied in two equal splits, half at the time of planting, and the remaining half 50 days after planting. Standard irrigation, weeding, and other management practices were followed when required throughout the growth period. Basil plants were harvested from the central part of each plot on 65 days after planting at full bloom. At the end of growth period (beginning of anthesis) three middle rows of each plot has been taken after eliminating the edge effects. Some agronomical parameters like plants height, stems diameter, inflorescence number and length, leaf area index, chlorophyll a+b, chlorophyll a/b ratio, chlorophyll content (SPAD value) and dry matter yield have been determined. Leaf surface of each plant determined using leaf area meter (USA LICOR Model LI-3000). Total leaf chlorophyll contentwas recorded using chlorophyll meter (SPAD – 502, Minolta, Japan). For measuring the amount of chlorophyll a and b (Equation 1-3) with methanol and measuring the absorbed light spectrum from the solution by spectrophotometer (6405.UV/Vis – Jenway - England) in 652 and 665.2 wavelengths (Porra et al., 1989). $[Chl a] = 16.29 E^{665.2} - 8.45 E^{652}$

 $[Chl b] = 30.66 E^{652} - 13.58 E^{665.2}$ $[Chl s a+b] = 22.12 E^{652} + 2.71 E^{665.2}$

Statistical analysis of data and orthogonal comparisons were done using SAS (SAS Institute, 1997) software and means were compared using least significant difference (LSD) test ($P \le 0.05$).

Propert ies	Electrical conductivit	pН	Organic matter	N	Р	K	Sand	Clay	Silt	Soil Texture
105	y (dS.m ⁻¹)		(%)		mg kg ⁻¹		(%)			
(Soil)	1.52	7.81	2.43	0.26	10	270	10.33	46.33	43.33	Silty-Clay
Sewage Sludge	10.75	7.44	4.50	0.58	43	480	-	-	-	-


Table 1. Chemical properties of soil and vermicompost in this study

Results

Morphological trait

Variance analysis of data showed that the type of fertilizer significantly affects plants height, stems diameter, inflorescence number and lenght, leaf area index and dry matter yield (Table 2).

Results represented that in $SS_{20}+50\%$ CF treatment, plants height significantly increased (11% and 6%, respectively) compared to both control and chemical fertilizer treatments (Fig 1: a). In addition, in chemical fertilizer treatment and SS₂₀+50% CF treatment highest stem diameter (9.42 and 9.66mm, respectively) was obtained (Fig 1: b). On the other hand, increasing the amount of sewage sludge increase (from SS_{20} to SS_{40} +50% CF), significantly decreased plants height and stems diameter by 11% and 34% respectively (Fig 1: a and b). According to results of this research, maximum inflorescence number and length (43 and 39 cm, respectively) and dry matter yield (5947 kg ha⁻¹) were recorded when SS_{20} treatment was applied. Application of this treatment (SS_{20}) increased the inflorescence number and length by 2-3 times and 41% compared to control, respectively (Fig 1: c and d). Furthermore, over application of sewage sludge (both SS_{40} and SS_{40} +50% CF treatments) markedly reduced inflorescence number and dry matter yield up to 50% and 54%, respectively. Meanwhile, the lowest leaf area index (LAI) was observed in control and SS₄₀+50% CF (Fig 1: e and f).



International Journal of Agricultural Technology 2013, Vol. 9(2): 451-460

Fig. 1. Effects of different amounts of fertilizers on morphological traits in basil CF: chemical fertilizer; SS20 : sewage sludge 20 Mg ha-1; SS20+½CF : sewage sludge 20 Mg ha-1+ 50% Chemical fertilizer; SS40: sewage sludge 40 Mg ha-1; SS40+½CF : sewage sludge 40 Mg ha-1 + 50% Chemical fertilizer.

Physiological traits

Results showed that sewage sludge, significantly affects the amount of chlorophyll a, chlorophyll b, chlorophyll a+b, chlorophyll a/b ratio and leaf chlorophyll content (SPAD) (Table 2).

The highest amount of chlorophyll a (4.88 mg ml⁻¹) observed in control (Table 2). Results in table 4 also showed that chlorophyll a has the highest positive and significant correlation with chlorophyll a+b ($r = 0.97^{**}$) and a negative and significant correlation with LAI ($r = -0.62^{**}$). There was a weak negative linear regression (Fig 2) between chlorophyll a and leaf area index during flowering stage. On the other hand the highest amount of chlorophyll b (2.14 mg ml⁻¹) obtained by application of SS₂₀+50%CF treatment (Table 2).

Results also showed that highest amount of chlorophyll a+b recorded in control and $SS_{20}+50\%$ CF treatment (6.78 & 6.86 mg ml⁻¹, respectively) whereas increasing amount of sewage sludge up to SS_{40} and $SS_{40}+50\%$ CF dramatically decreased chlorophyll a+b content by 17% & 8%, respectively. Highest chlorophyll a/b ratio obtained by applying $SS_{40}+50\%$ CF treatment (Table 2). Results suggested that leaf chlorophyll content increased in chemical fertilizer and sewage sludge treatments, as the highest leaf chlorophyll content (48.07) was observed in SS_{20} treatment (Table 2).

	(Chlorophy	7 11	Chlorophyll	SPAD	
Treatment	а	b	a+b	a/b ratio	Value	
	(mg/ml)					
Control	4.88 ^a	1.90 ^b	6.78 ^a	2.57 °	35.90 ^d	
CF	3.36 ^f	1.36 ^e	4.71 ^d	2.48^{d}	46.85 ^{ab}	
SS20	4.02 ^e	1.61 ^c	5.63 °	2.50^{d}	48.07^{a}	
SS20+1/2 CF	4.72 °	2.14 ^a	6.86 ^a	2.21 ^e	41.25 °	
SS40	4.17 ^d	1.51 ^d	5.68 °	2.76 ^b	43.20 bc	
SS40+1/2 CF	4.76 ^b	1.56 °	6.32 ^b	3.04 ^a	45.80^{ab}	
F test						
Mean Squares (MS)	1.03**	0.25**	2.01**	0.24**	60.17**	
Coefficient of Variation (CV)	0.4	1.7	0.8	1.1	5.2	

Table 2. Effects of Different amounts of fertilizer treatments on basi leaf

 chlorophyll content

In each columns means with similar letters were not significantly different based on LSD test. **: significant at 1% levels

Table 3. Simple correlation between leaf area index and chlorophyll (n=18)

	Chlorophyll	Chlorophyll a	Chlorophyll	Leaf	area
	a+b		b	index	
Chlorophyll a+b	1				
Chlorophyll a	0.97**	1			
Chlorophyll b	0.87**	0.73**	1		
Leaf area index	-0.47*	-0.62**	0.09^{ns}	1	
ns, * and ** non- s	ignificant and sign	ificant at 5 and 1%	levels, respective	ly	

International Journal of Agricultural Technology 2013, Vol. 9(2): 451-460




Fig. 2. Regression relationship between chlorophyll a and leaf area index (LAI)

Discussion

The significant effects of fertilizers type in different amounts on morphological traits (plants height, stems diameter, inflorescence number and length, leaf area index and dry matter yield) and leaf chlorophyll amounts (chlorophyll a, chlorophyll b, chlorophyll a+b, chlorophyll a/b ratio and leaf chlorophyll content (SPAD) showed that, sewage sludge, which is a kind of organic waste, increases macro and micro elements like nitrogen, phosphorus, potassium, iron, manganese, cooper and zinc (Baran et al., 2001). Meanwhile, using it plus chemical fertilizers, improves physical characteristics of soil, and increases the production by using water and nutrients efficiently (Bhattacharyya, 2008). Thus by applying sewage sludge, nitrogen absorption increases in roots which raises the growth and production of more leaves and inflorescences (Sharifi et al., 2010) and therefore, causes an increase in dry matter yield. Delgado et al. (2003) reported that adding 1000 kg sewage sludge plus 350 kg urea per hectare, increases the corns yield by 20% compared to control, Shabanian Borujeni *et al.* (2005) also noticed that 50 Mg ha⁻¹ and 100 mg ha⁻¹ of sewage sludge treatments causes a significant increase in aboveground dry matter weight and plant height in snapdragon (Antirrhinum majus) compared to control. This researchers also found that number of florets in snapdragon inflorescences also increased from 13 in control treatment to around 19 in 50 Mg ha⁻¹ sewage sludge treatment. They suggested that presence of organic matter and nutrients like nitrogen, phosphorus, potassium, iron and other nutrients in sewage sludge caused the increase in snapdragon growth. Similarly, Perez-Murcia et al. (2006) found that the maximum fresh and dry weight of Broccoli (Brassica oleracea) obtained in a 30% sewage sludge combination. They suggested that adding an appropriate amount of sewage sludge to soil can supply the plants with nutrients in a better situation. In Spinach (Spinacia oleracea) Vaseghi et al. (2005) also observed an increase in the dry and fresh matter yield along with increase of sewage sludge application. They believed that the main reason is the increase of nitrogen absorption by roots and production of more leaves. Higher light absorption and photosynthesis following the increase in number of leaves lead to production of more

hydrocarbons in leaves and caused the increase in number of blossoms and dry matter yield. Consistently, Gardio *et al.* (2005) noticed that application of sewage sludge caused the faba bean plant to be more green and have higher leaf surface and dry matter yield compared to control. Findings by Sabye *et al.* (1990) sage (*Artemisia* spp) and Atriplex (*Atriplex* spp) showed that leaf area index were significantly higher in the sections in which sewage sludge was applied; compared to control sections with no sewage sludge. Shabanian Borujeni *et al.* (2005) found that in clove gilly flower (*Dianthus spp*) chlorophyll density increased from 478 mg kg⁻¹ in control treatment to 596 mg kg⁻¹ and 726 mg kg⁻¹ in 50 Mg ha-1 and 100 Mg ha⁻¹ sewage sludge treatments, respectively. According to the low leaf area index, they believed that this increase in chlorophyll is because of increased chlorophyll production (high magnesium density in sewage sludge). Because magnesium is one of the main elements in chlorophyll structure.

Organic matter in sewage sludge improves soil characteristics including hydraulic conductivity, soil particles stability, aeration and moisture and thus improves growth and increases crops yield (Debosz et al., 2002) but on the other hand, inordinate application, increases the probability of toxic densities of elements and causes aggregation of heavy elements in soil (Amini *et al.*, 2008). Shirani et al. (2010) found that over application of sewage sludge causes excessive salinity of soil and decreases crops yield. Kassray and Saedi (2010) also noticed that application of 5 and 10 Mg ha⁻¹ sewage sludge treatments increased the fresh and dry weight of aboveground in tomato compared to control. Application of higher levels of sewage sludge a severe increase in yield and plant toxicities; as this decrease in 30 Mg ha⁻¹ level was estimated around 20% for fresh weight and 44% for dry weight compared to control. Parida and Das (2005) also found that one reasons of chlorophyll reduction in high salinity conditions is increased amount of chlorophyllase that degrades the chlorophyll. High salinity prevents the composition of chlorophyll and carotenoids in many plant species leaves and also induces senescence, chloroplast infraction and reduction of leaves chlorophyll in plants and this reduction causes less efficiency in photosynthesis.

Consistently, in the present study results showed that in higher amounts of sewage sludge application either enriched or none-enriched with chemical fertilizers (SS_{40} or $SS_{40}+50\%$ CF treatments) most of the studied traits (plants height, stems diameter, inflorescence number and length, leaf area index, dry matter yield, chlorophyll b, and chlorophyll a+b) were significantly reduced. This reduction is the consequence of excessive soil salinity due to effects of residuals of some elements particularly chloride, because in this research fertilizer treatments added to soil in four continuous years (2006-2009) and also effects of remaining elements during four years increased the probability of toxic densities of elements in soil or caused higher salinity and interruption in nutrients balance and finally reduced the plants growth. Hence application of SS_{20} and $SS_{20}+50\%$ CF treatments could cause better improvement in the growth and increase of the yield in basil compared to SS_{40} and $SS_{40}+50\%$ CF treatments.

Conclusion

Generally it is observed that 20 Mg ha⁻¹ of sewage sludge enriched or none-enriched with chemical fertilizer treatments have the best effect on studied traits compared to control and chemical fertilizer treatment. As application of this treatment caused an increase in some traits (plants height, inflorescences number and length, leaf area index, dry matter yield and chlorophyll a+b) in basil leaves compared to control and chemical fertilizer treatment. Thus application of this fertilizer (sewage sludge) instead of a part of chemical fertilizer in basil cultivation, will reduce fertilizer consumption and optimize the usage of sewage sludge in addition to promotion of morphological traits and chlorophyll content.

References

- Amini, H, Ansari, A. (2008). Sewage impact on heavy metal uptake by spinach. Proceedings of the Third National Congress of Recycling and Using Renewable Organic Resources in Agriculture. In Isfahan.
- Shabanian Borojeni, H., Haj Abasi, M.A., Mobli, M. and Afuni, M. (2005). Effects of waste water and sewage sludge of Iran polyacrile factory on some morphological characteristics and ION concentration of lawn, snapdragon and sweet williams. Iranian Journal of Horticultural Science 6(3):135-148.
- Akbarnejad, F., Astaraee, A.R., Fotovat, M. and Nasiri Mahalati, M. (2010). Effects of municipal compost and sewage sludge on yield and yield components of black cumin (*Nigella sativa L.*) medicinal plants. Iranian Journal of Field Crops Research 8(5):767-771.
- Baran, A., Cayci, G., Kutuk, C. and Hartmann, R. (2001). The effect of grape marc as growing medium on growth of hypostases plant. Bioresource Technology 78:103-106.
- Bhattacharyya, R., Kundu, S., Prakash, V. and Gupta, H.S. (2008). Sustainability under combined application of mineral and organic fertilizers in a rain fed soybean-wheat system of the Indian Himalayas. European Journal of Agronomy 28:33-46.
- Darzi, M.T., Ghalavand, A. and Rejali, F. (2008). Effect of mycorrhiza, vermicompost and phosphate biofertilizer application on flowering, biological yield and root colonization in fennel (*Foeniculum vulgare* Mill.). Iranian Journal of Crop Science 10(1):88-109.
- Debosz, K., Petersen, S.O., Kure, L.K. and Ambus, P. (2002). Evaluating effects of sewage sludge and household compost on soil physical, chemical and microbiological properties. Applied Soil Ecology 19(3):248-237.
- Delgado, M.M., Parcel, M.A., Miralles, R., Beltrain, E.M., Bernigole, L. and Valero, J. (2003). Management sewage sludge thermal drying amended soil. Waste Contaminants: Lifecycle and Entry into food chain. pp. 89-92.
- Ferreras, I., Gomez, E., Toresani, S., Firpo, I. and Rotondo, V. (2006). Effect of organic amendments on some physical, chemical and biological properties in a horticultural soil. Bioresource Technology 97:635-640.
- Gardio, S., Martin del champion, G., Esteller, M.V., Vaca, R. and Lugedo, J. (2005). Heavy metals in soil treated with sewage sludge composting, their effect on yield and uptake of broad bean seeds (*Vicia faba* L.). water Air and Soil Pollution 166:303-319.
- Haghighi, M. (2011). Sewage Sludge Application in Soil Improved Leafy Vegetable Growth. J Biological Environmen and Sciences 5(15):165-167.

- Kassray, R. and Saedi, S. (2010). Effects of Tabriz petrochemical sewage sludge on tomato growth. Journal of Water and Soil. 24(1):10-20.
- Magdoff, F. and Weil, R.R. (2004). Soil organic matter in sustainable agriculture. CRC Press, Boca Raton. pp. 398.
- Parida, A.K. and Das, A.B. (2005). Salt tolerance and salinity effects on plant: A review. Ecotoxicology and Environmental Safety 60:324-349.
- Perez-Murcia, M.D., Moral, R., Moreno-Caselles, J. and Paredes, A.C. (2006). Use of composted sewage sludge in growth media for broccoli. Bioresource Technology 97:123-130.
- Porra, R.J., Thompson, W.A. and Kriedemann, P.E. (1989). Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectrometry. Photosynthesis Research 975:384-394.
- Sabye, B.R., Rendelton, R.L. and Webb, R.L. (1990). Effect of municipal sewage sludge application on two reclamation shrub species in copper mine spoils. Journal of Environmental Quality 19:580-586.
- SAS Institute. Inc. (1997). SAS/STAT Users Guide, Version 6.12. SAS Institute. Inc. Cary, NC.
- Sharifi, M., Afyouni, M. and Khoshgoftarmanesh, A.H. (2010). Effects of Animal Manure, Sewage Sludge, and Cadmium Chloride on Cadmium Uptake of Corn Shoots. Iranian Journal of Water and Sewage 4:98-103.
- Shirani, H., Hajabbasi, M.A., Afyouni, M. and Dashti, H. (2010). Cumulative Effects of Sewage Sludge on Soil Physical and Chemical Characteristics. Iranian Journal of Water and Sewage 3:28-36.
- Shober, A.L., Stehouwer, R.C. and Macneal, K.E. (2003). On-farm assessment of biosolids effects on soil and crop tissue quality. Journal of Environmental Quality 32:1873-1880.
- Singh, R.P. and Agrawal, M. (2008). Potential benefits and risks of land application of sewage sludge. Waste Management Research 28(2):347-358.
- Spnmez, F. and Bozkurt, M.A. (2006). Lettuce grown on calcareous soils benefit from sewage sludge. Journal of Acta Agriculture Scandinavica 56(1):17-24.
- Vaseghi, S., Afyouni, M., Shariat Madari, H. and Mobli, M. (2005). Effect of sewage sludge on same macronutrients concentration and soil chemical properties. Journal of Water and Wastewater 53:18-25 (In Persian).
- Warman, P.R. and Termeer, W.C. (2005). Evaluation of sewag sludge, septic waste and sludge compost applications to corn and forage: Ca, Mg, S, Fe, Mn, Cu, Zn and B content of crops and soils. Bioresource Technology 96:1029-1038.

(Received 16 October 2012; accepted 28 Febuary 2013)