
Repellant activity of plant essential oils formulation against three disease causing mosquito vectors

A. Manimaran¹, M. Mary Jee Jee Cruz¹, C. Muthu^{2*}, S. Vincent³ and S. Ignacimuthu²

¹Department of Biosciences, AVIT, Vinayaka Missions University, Paiyanoor, Chennai - 603 104, India, ²Entomology Research Institute, Loyola College, Chennai 600 034, ³Department of Advanced Zoology and Biotechnology, Loyola College, Chennai 600 034, India.

A. Manimaran, M. Mary Jee Jee Cruz, C. Muthu, S. Vincent and S. Ignacimuthu (2013). Repellant activity of plant essential oils formulation against three disease causing mosquito vectors. International Journal of Agricultural Technology 9(4):845-854.

Abstract A plant oil formulation (POF) consisting of calamus, cinnamon, citronella, clove, eucalyptus, lemon, menthe, orange oils and natural Camphor 10% was evaluated for its repellant potential against *Aedes aegypti* (L.), *Anopheles stephensi* Liston and *Culex quinquefasciatus* Say at 250 and 15.25 ppm. At 250 ppm concentrations the POF provided 100% protection from bites of *Anopheles stephensi* up to a mean time of 6 hours and 30 minutes while for *C. quinquefasciatus* and *A. aegypti*, it was 5 h 30 min. At 15.62 ppm it also exhibited more than 74% mean protection against *C. quinquefasciatus* and *A. aegypti* while it was 62.88% effective against *A. stephensi*. Ethanol which was used for suspension was kept as negative control and DEET (25 mg) served as positive control. The study concludes that this formulation could be used to repel these insects.

Key words: Biting protection, DEET, *Aedes aegypti*, *Anopheles stephensi*, *Culex quinquefasciatus*

Introduction

Mosquitoes act as vectors for the transmission of dangerous diseases like malaria, lymphatic filariasis and viral diseases which lead to morbidity, mortality, social and economical loss (Becker *et al.*, 2003). Personal protective measures, including repellants are widely used to prevent the transmission of arthropod-borne diseases by minimizing the contact between humans and vectors. In contrast to vaccine and chemoprophylaxis as means of personal protection, repellants are considered as convenient, inexpensive and affordable with advantages in protection against a wide range of vectors (WHO, 1995). Different volatile organic molecules have the potential to inhibit the scent

* Corresponding author: C. Muthu; e-mail: muthu_mathi@yahoo.com

orientation of insects in confined environmental conditions and therefore can be efficiently used as repellants against mosquito populations (Oyedele *et al.*, 2002; Kline *et al.*, 2003; Achee *et al.*, 2009; Drapeau *et al.*, 2009). Repellancy is one of the commonly used methods to prevent mosquitoes and other blood-sucking pests from biting. Repellants are applied directly on the skin or to clothing and other fabrics such as bed nets and anti-mosquito screens. Repellants evaporate much more quickly than most insecticides. The prolonged use of commercially developed mosquito coils, mats and vapourizers which contains synthetic pyrethroids like pyrethrin, allethrin and paraallethrin and chemical such as DEET may cause harmful effects (Bell *et al.*, 2002); also, the mosquitoes develop resistance to these chemicals (Magurayni *et al.*, 2009). In the commercial mosquito vapourizers, the synthetic and chemical insecticides are dissolved in deodourized kerosene, which also causes side effects to the user. In search of alternative and safe methods to protect from mosquito bites, products from essential oils are emerging as good mosquito repellants and they are effective, safe and also low cost (Mittal *et al.*, 2003; Senthil Nathan *et al.*, 2004).

Researchers all over the world have reported the repellent activity of essential oils derived from plants against the adults of various mosquito species. Volatile oils from *Acorus calamus* showed repellent activity against *A. Aegypti* (Tariq and Quadri, 2001); oil from *Cinnamomum camphora*, *C. zeylanicum*, and *Citrus limon* (Amer and Mehlhorn, 2006), *C. zeylanicum* (Prajapati *et al.*, 2005) form chemical repellants against *A. aegypti*, *An. stephensi* and *C. quinquefasciatus*. Oil from *Citronella spp.* has shown repellent activity against *A. aegypti* (Nazli *et al.*, 2008).

Essential oils from leaves of *Cymbopogon nardus* and *Cymbopogon sp.* showed repellent activity against *A. aegypti*, *A. stephensi*, *A. culicifacies* and *C. quinquefasciatus* (Ansari and Razdan, 1995; Tyagi *et al.*, 1998; Trongtokit *et al.*, 2005); *Eucalyptus globulus* oil revealed repellent activity against *A. aegypti* (Choi *et al.*, 2002), *A. albopictus* (Yang and Ma, 2005), *C. pipiens pallens*, *A. stephensi* and *C. quinquefasciatus* (Amer and Mehlhorn, 2006), *Mentha piperita* oil exhibited repellent activity against *A. stephensi*, *A. aegypti* and *C. quinquefasciatus* (Pathak *et al.*, 2000; Amer and Mehlhorn, 2006) while oil from *M. spicata* and *Citrus sinensis* showed repellency against *A. albopictus* (Yang and Ma, 2005) and *C. pipiens molestus* (Traboulsi *et al.*, 2005). Rajkumar and Jebanesan (2007) showed the repellent activity of essential oils from *Centella asiatica*, *Ipomoea cairica*, *Momordica charantia*, *Psidium guajava* and *Tridax procumbens* against the malarial vector, *A. stephensi*.

Combination of different oils such as *Syzygium* spp., *Cymbopogon* spp., *Eucalyptus* spp., *Pelargonium* spp., *Lavandula* spp., *Mentha* spp., *Santalum* spp. and *Thymus* spp. exhibited repellent activity against *A. aegypti* (Girgenti and Suss, 2002). Recently, essential oil formulations have been reported to have repellent activity against *A. albopictus* and *C. quinquefasciatus* (Cliek *et al.*, 2010), *A. gambiae* and *Anopheles funestus* (Oparaocha *et al.*, 2010). Lawal *et al.* (2012) reported that different formulations from six essential oils showed repellent activity against *A. gambiae*. The present study was aimed at evaluating the repellent activity of an essential oil formulation comprising eight oils against *A. aegypti*, *A. stephensi* and *C. Quinquefasciatus*.

Materials and methods

Based on preliminary screening (Manimaran *et al.*, 2012), the following oils were selected for this study. The plant oil formulation (POF) consisted of calamus (*Acorus calamus* Linn.) oil – 3%, cinnamon (*Cinnamomum veerum* Presl.) oil – 13%, citronella (*Cymbopogon nardus* Rendle) oil – 12%, clove (*Myrtus caryophyllus* L.) oil – 24%, eucalyptus (*Eucalyptus globulus* L.) oil – 6%, lemon (*Citrus limon* (L.) Burm. F.) oil – 7%, mentha (*Mentha piperita* L.) oil – 10%, orange (*Citrus sinensis* (L.) Osbeck) oil – 25% and natural Camphor 10% (10 g in 100 mL).

The repellency of volatile oils was evaluated using the human-bait technique (Schreck and McGovern, 1989; WHO, 2009). The repellent activity was determined by the percentage of protection in relation to dose method (WHO, 1996). Three to four days old, blood starved 100 adult female *Anopheles stephensi*, *Culex quinquefasciatus* and *Aedes aegypti* mosquitoes were kept in respective net cages (45 cm x 30 cm x 45 cm). The forearms of the test persons were cleaned with distilled water. After air drying, one arm of the test person (only 2.5 cm² dorsal side of the skin) was exposed and the remaining area being covered by rubber gloves. The POF was dissolved in ethanol to make 15.25 and 250 ppm concentrations. About 1 mL of the POF was applied to the exposed part of the forearm of volunteers. One volunteer applied with ethanol served as negative control. DEET (25 mg) was used as positive control.

The periodicity of feeding pattern differed from species to species. According to the feeding time, the repellency tests were conducted. At each testing period the hands were introduced inside the test cage and exposed to the adult mosquitoes for 5 min duration from 6:00 h until 18:00 h for *Aedes aegypti* and *Anopheles stephensi* and from 20:00 h until 06:00 h for *Culex quinquefasciatus*. Each test was replicated six times. The number of mosquitoes

landing on both the control and treated hands was recorded at every 5 minutes exposure period with the help of a timer.

$$\text{Percentage of mosquitoes fed} = \frac{\text{No. of mosquitoes fed}}{\text{Total no. of mosquitoes released}} \times 100$$

$$\text{Percentage of protection} = \text{Percentage of mosquitoes fed} - 100$$

Results

The results of repellent activity of POF on the forearm of the volunteers in the laboratory against *Anopheles stephensi* are presented in Table 1. At 250 ppm, the POF provided 100% protection from bites of *Anopheles stephensi* up to a mean time of 6 h 30 min. The control gave protection for only about 30 min. The study clearly showed that the application of 250 ppm concentration of the POF provided 83.19 % biting protection from *Anopheles stephensi*. DEET (25 mg) recorded the biting protection for 5.30 h and it showed 80.34% biting protection.

Table 1. Repellent Activity of the Plant Oil Formulation Against the Adult *Anopheles stephensi*

Observation Time	Mean number of mosquitoes Landing / Feeding on volunteers			
	Plant Oil (250 ppm)	Plant Oil (15.62 ppm)	Ethanol (Control)	DEET (25 mg)
18:00 – 19:00	0	0	0	0
19:01 – 20:00	0	0	6.83 ± 0.40	0
20:01 – 21:00	0	3.33 ± 0.51	8.33 ± 1.03	0
21:01 – 22:00	0	4.33 ± 0.51	7.66 ± 1.03	0
22:01 – 23:00	0	4.16 ± 0.40	6.33 ± 0.51	0
23:01 – 24:00	0	3.16 ± 0.40	8.33 ± 0.51	2.17±0.41
00:01 – 01:00	4.00 ± 0.63	5.00 ± 0.63	8.16 ± 0.75	2.50 ±0.55
01:01 – 02:00	3.16 ± 0.40	3.66 ± 0.81	7.33 ± 1.03	2.67 ± 0.82
02:01 – 03:00	3.16 ± 0.51	4.83 ± 0.75	4.16 ± 0.75	3.17±0.75
03:01 – 04:00	2.00 ± 0.63	3.16 ± 0.81	6.83 ±0.40	3.66±0.81
04:01 – 05:00	2.16 ± 0.40	3.16 ± 0.75	7.66 ± 1.03	3.16±0.40
05:01 – 06:00	2.33 ± 0.51	2.33 ± 0.51	4.50 ± 0.83	2.33±0.51
Total (% mosquito fed)	16.81±1.58	37.12 ± 1.03	76.12 ± 0.98	19.66±1.50
% of Protection	83.19	62.88	23.88	80.34
Mean protection time (Min)	390.00	150	30	330.00

Values represent Mean ± Standard deviation of six replicates.

*Value represent Mean ± SD of the means

The repellent activity of the POF against *Culex quinquefasciatus* is given in Table 2. At 250 ppm 100% protection from the bites of *Culex quinquefasciatus* up to a mean time of 5 h 30 min was observed. The control protected only 30 min from mosquito bite or landing. The POF at 250 ppm provided 87.19% biting protection from *Culex quinquefasciatus*. DEET (25 mg) recorded the biting protection for 5.30 h and it showed 81.53% biting protection.

Table 2. Repellent Activity of the Plant Oil Formulation Against the Adult *Culex quinquefasciatus*

Observation Time	Mean number of mosquitoes Landing / Feeding on volunteers			
	Plant (250 ppm)	Oil Plant (15.62 ppm)	Oil	Ethanol (Control)
18:00 – 19:00	0	0	0	0
19:01 – 20:00	0	0	5.16 ± 0.75	0
20:01 – 21:00	0	0	7.16 ± 0.75	0
21:01 – 22:00	0	2.33 ± 0.51	8.16 ± 0.75	0
22:01 – 23:00	0	1.16 ± 0.40	7.33 ± 1.03	0
23:01 – 24:00	3.00 ± 0.63	2.00 ± 0.63	8.33 ± 1.03	1.833±0.41
00:01 – 01:00	2.16 ± 0.40	3.50 ± 0.83	9.66 ± 0.75	2.00 ± 0.63
01:01 – 02:00	2.33 ± 0.51	2.16 ± 0.40	7.33 ± 1.03	2.33 ± 0.52
02:01 – 03:00	1.16 ± 0.40	4.33 ± 0.51	4.50 ± 0.83	2.66±0.52
03:01 – 04:00	2.00 ± 0.63	2.83 ± 0.40	8.16 ± 0.75	2.83±0.75
04:01 – 05:00	1.16 ± 0.40	2.00 ± 0.63	6.33 ± 0.51	3.16±0.40
05:01 – 06:00	1.00 ± 0.63	3.16 ± 0.40	5.50 ± 0.83	3.66±0.82
Total (mosquito fed %)*	12.81 ± 0.83	23.47 ± 1.03	75.62 ± 0.98	18.47±1.44
% Protection	87.19	76.53	24.38	81.53
Mean protection time (Min)	330	180	60	330.00

Values represent Mean ± Standard deviation of six replicates.

*Value represent Mean ± SD of the means

At 250 ppm of POF, it provided 100% protection from bites of *Aedes aegypti* up to a mean time of 5 h 30 min. The control recorded only 30 min protection from mosquito bite or landing. The POF at 250 ppm concentration provided 87.03 % biting protection from *Aedes aegypti* (Table 3). DEET (25 mg) recorded the biting protection for 5.30 h and it showed 84.33% biting protection.

Allergic reaction study clearly indicated that the POF did not show any allergic effects such as itching, breathing difficulty, or headache to the volunteers.

Table 3. Repellant Activity of the Plant Oil Formulation Against the Adult *Aedes aegypti*

Observation Time	Number of Mosquitoes Landing / Feeding on Volunteers					
	Plant (250 ppm)	Oil	Plant (15.62 ppm)	Oil	Ethanol (Control)	DEET (25 mg)
06:00 – 07:00	0		0		0	0
07:01 – 08:00	0		0		7.33 ± 1.03	0
08:01 – 09:00	0		0		9.16 ± 0.75	0
09:01 – 10:00	0		2.16 ± 0.75		8.33 ± 1.03	0
10:01 – 11:00	0		1.33 ± 0.51		6.83 ± 0.40	0
11:01 – 12:00	2.33 ± 0.51		2.33 ± 0.51		5.16 ± 0.75	1.5±0.57
12:01 – 13:00	3.33 ± 0.51		5.00 ± 0.63		8.33 ± 1.03	1.67±0.52
13:01 – 14:00	1.50 ± 0.63		3.33 ± 0.51		4.50 ± 0.83	2.00±0.63
14:01 – 15:00	2.16 ± 0.40		3.50 ± 0.83		6.83 ± 0.40	2.33±0.52
15:01 – 16:00	1.33± 0.51		2.16 ± 0.75		7.66 ± 1.03	2.67±0.52
16:01 – 17:00	1.16 ± 0.40		2.16 ± 0.75		6.16 ± 0.75	3.0±0.63
17:01 – 18:00	1.16 ± 0.40		3.33 ± 0.51		5.15 ± 0.75	2.5±0.55
Total (Mosquito fed %)	12.97 ± 1.58		25.14 ± 1.03		75.45 ± 0.98	15.67±1.22
% of Protection Mean	87.03		74.86		24.55	84.33
protection time (Min)	330		180		30	330.00

Values represent Mean ± Standard deviation of six replicates.

*Value represent Mean ± SD of the means

Discussion

The present study clearly revealed that the formulation at 250 ppm offered a mean protection time of more than 330 minutes (i.e. 5h. 30 min) against the mosquito species tested. The result of this study coincides with the earlier results of Das *et al.* (2003) who reported that *Zanthoxylum limonella* and *Citrus aurantifolia* essential oils when mixed with mustard oil exhibited a mean protection time of more than 302 min against *Aedes (S.) albopictus* at 30% concentration. Complex mixture of five essential oils recorded more than 450 minutes protection time against *A. aegypti*, *C. Quinquefasciatus* and *A. stephensi* (Amer and Mehlhorn, 2006). Khandagle *et al.* (2011) reported that essential oil from rhizome of *Zingiber officinalis* recorded 300 min protection from biting of *A. aegypti* and *C. quinquefasciatus*. An oil formulation containing *Nepeta cataria* was tested against black flies, *Simulium decorum* and mosquito *Aedes intrudens* for its repellant activity. The formulation exhibited over 4 hrs

protection from bites (Spero *et al.*, 2008). Oyedele *et al.* (2002) reported that the products (ointment and cream) from lemon grass oil exhibited 2-3 hrs protection from biting of *A. aegypti* at a concentration of 15%v/w. In the present investigation, the concentration of 250 ppm of the POF exhibited 83.19% protection against *Anopheles stephensi*. The concentration of 250 ppm of the POF exhibited 87.19 % protection against *Culex quinquefasciatus*. The POF at 250 ppm concentration exhibited 87.03% protection against *Aedes aegypti* bite. Amer and Mehlhorn (2006) reported that a mixture of essential oils provided biting protection against *A. aegypti*, *C. Quinquefasciatus* and *A. stephensi*. The essential oil formulation comprising of *E. globulus*, *O. basilicum*, *C. citratus*, *C. sinensis*, *A. indica* (Neem) and *H. suaveolens* exhibited 72.22% protection (2 h) from biting of *A. gambiae* at 10% concentration (Lawal *et al.*, 2012). *E. globulus* and *Ocimum basilicum* mixed in soybean oil exhibited 99% biting protection from *A. aegypti* and *A. dirus* with protection time of 210 minutes (Sritabutra *et al.*, 2011). The repellent activity in the present study may be due to the presence of *E. globulus*, *C. citrates* and *C. sinensis* which were used in the formulation along with other essential oils. Similarly Trongtokit *et al.* (2005) reported that the essential oils from *C. nardus*, *Pogostemon cablin*, *Syzygium aromaticum* and *Zanthoxylum limonella* effectively provided 100% repellency for 2 hrs from biting of *A. aegypti*. Combination of different oils such as *Syzygium* spp., *Cymbopogon* spp., *Eucalyptus* spp., *Pelargonium* spp., *Lavandula* spp., *Mentha* spp., *Santalum* spp., and *Thymus* spp. exhibited repellent activity against *Ae. Aegypti* (Girgenti and Suss, 2002). The repellent activity of plant products may be due to the active principles present in the plants which block mosquito sensory organs with heavy, bulky molecules (Ned Roze, 2012).

In the present investigation the essential oil formulation showed repellent activity against *Anopheles stephensi*, *Culex quinquefasciatus* and *Aedes aegypti*. In this study the repellent activity depends on the concentration of POF (250 ppm and 15.62 ppm). The present findings corroborate with the findings of Lawal *et al.* (2012) who reported that the percentage protection was concentration dependent manner. Higher concentration showed higher protection than the lower concentration. Similarly, Venkatachalam and Jebanesan (2001) reported that *Feronia elephantum* exhibited the total percentage protection of 45.8 % at 1.0 mg/cm² and 59.0 % at 2.5 mg/cm². Essential oil formulation comprising of *Ocimum gratissimum*, with olive oil at 30% concentration showed more than 96% repellency while 20% concentration recorded less than 75% repellency against *A. gambiae* and *Anopheles funestus* (Oparaocha *et al.*, 2010).

The POF contains some chemical compounds like eugenol in clove oil, 1,8-cineole in *Eucalyptus globules*, citronellal in lemon grass, menthol and pulegone in menthe oil, cinnamaldehyde, eugenol and cinnamyl acetate in cinnamon oil, several terpenes and geraniol in lemon oil. These maybe responsible for repellent activity of five essential oils containing monoterpenes such as eucalyptol, geraniol, limonene, linalool, menthone, linalyl acetate and menthyl acetate which has been shown effective against blood sucking bug, *Rhodnius prolixus* which is a vector of Chagas disease in many Latin American nations (Sfara *et al.*, 2009). Choi *et al.* (2002) reported that the presence of α -terpinene, carvacrol and thymol in *Lavandula officinalis* and *R. officinalis* were responsible for the repellent activity against *C. pipiens pallens* adults.

Conclusion

The plant oil formulation exhibited repellent activity against the tested vector mosquitoes more than 300 minutes. Use of these botanical derivatives in mosquito control instead of synthetic insecticides could reduce the cost and has no adverse environment effects.

Acknowledgements

The authors would like to thank Entomology Research Institute, Loyola College, Chennai for financial support for carried out this work.

References

- Achee, N.L., Sardelis, M.R., Dufour, I., Chauhan, K.R. and Grieco, J.P. (2009). Characterization of spatial repellent, contact irritant, and toxicant chemical actions of standard vector control compounds. *Journal of the American Mosquito Control Association* 25:156-167.
- Amer, A. and Mehlhorn, H. (2006). Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitology Research* 99: 466–472.
- Ansari, M.A. and Razdan, R.K. (1995). Relative efficacy of various oils in repelling mosquitoes. *Indian journal of malariology* 32:104-111.
- Becker, N., Petrie, D., Zgomba, M., Boase, C., Dahl, C., Lane, J. and Kaiser, A. (2003). *Mosquitoes and their control*. New York: Kluwer Academic/Plenum Publishers.
- Bell, J.W., Veltri, J.C. and Page, B.C. (2002). Human exposures to N,N-diethyl-*m*-toluamide insect repellents reported to the American Association of Poison Control Centers, 1993–1997. *International Journal of Toxicology* 2:341-52.
- Choi, W.S., Park, B.S., Ku, S.K. and Lee, S.K. (2002). Repellent activities of essential oils and monoterpenes against *Culex pipiens*. *Journal of the American Mosquito Control Association* 18:348–351.

- Cilek, J.E., Hallmon, C.F. and Johnson, R. (2010). Limited Efficacy of Commercially Formulated Essential Oils on Vegetation against Female *Aedes albopictus* and *Culex quinquefasciatus*. *Journal of Agricultural and Urban Entomology* 27:54-60.
- Das, N.G., EBaruah, I., Talukdar, P.K. and Das., S.C. (2003). Evaluation of botanicals as repellants against mosquitoes. *Journal of Vector Borne Diseases* 40:49-53.
- Drapeau, J., Frohler, C., Touraud, D., Krochel, U., Geier, M., Rose, A. and Kunz, W. (2009). Repellent studies with *Aedes aegypti* mosquitoes and human olfactory tests on 19 essential oils from Corsica, France. *Flavour and Fragrance J.* 24:160-169.
- Girgenti, P. and Suss, L. (2002). Repellent activity against *Aedes aegypti* (L) from formulas based on natural vegetable extracts are synthetic active agents *Annual Review of Immunology* 14:205-210.
- Khandagle, A.J., Tare, V.S. Raut, K.D. and Morey, R.A. (2011). Bioactivity of essential oils of *Zingiber officinalis* and *Achyranthes aspera* against mosquitoes. *Parasitology Research* 109:339-343.
- Kline, D.L., Bernier, U.R., Posey, K.H. and Branard, D.R. (2003). Olfactometric evaluation of spatial repellants for *Aedes aegypti*. *Journal of Medical Entomology* 40: 463-467.
- Lawal, H.O., Adequyi, G.O., Fawehinmi, A.B., Adeogun, A.O. and Etatuvie, S.O. (2012). Bioassay of herbal mosquito repellent formulated from the essential oil of plants. *J Journal of Natural Products* 5:109-115.
- Maguranyi, S.K., Webb, C.E., Mansfield, S. and Russel, R.C. (2009). Are commercially available essential oils from Australian native plants repellent to mosquitoes? *Journal of the American Mosquito Control Association* 25:292-300.
- Manimaran, A., Cruz, M., Muthu, C., Vincent, S. and Ignacimuthu, S. (2012). Larvicidal and knockdown effects of some essential oils against *Culex quinquefasciatus* Say, *Aedes aegypti* (L.) and *Anopheles stephensi* (Liston). *Advances in Bioscience and Biotechnology* 3:855-862.
- Mittal, P.K. and Subbarao, S.K. (2003). Prospect of using Herbal Product in the Control of Mosquito vectors. *ICMR Bulletin* 33: 0377-4910.
- Nazli, R., Akhter, M., Ambreen, S., Solangi, A.H. and Sultana, N. (2008). Insecticidal, nematocidal and antibacterial activities of *Gliricidia sepium*. *Pakistan Journal of Botany* 40:2625-2629.
- Ned Roze, 2012. *Snow Mosquitoes* the First Wave of Summer irritants, Alaska Science Forum, April 11, 2012, Article #2110.
- Oparaocha, E.T., Iwu, I. and Ahanakuc, J.E. (2010). Preliminary study on mosquito repellent and mosquitocidal activities of *Ocimum gratissimum* (L.) grown in eastern Nigeria. *Journal of Vector Borne Diseases* 47:45-50.
- Oyedela, A.O., Gbolade, A.A., Sosan, M.B., Adewoyin, F.B., Soyely, O.L. and Orafidiya, O.O. (2002). Formulation of an effective mosquito repellent topical product from lemon grass oil. *Phytomedicine* 9:259-262.
- Pathak, N., Mittal, P.K., Singh, O.P., Sagar, V. and Vasudevan, P. (2000). Larvicidal action of essential oils from plants against the vector mosquitoes *Anopheles stephensi* (Liston) *Culex quinquefasciatus* (Say) and *Aedes aegypti* (L). *International Pest Control* 42:53-55.
- Prajapati, V., Tripathi, A.K., Aggarwal, K.K. and Khanuja, S.P.S. (2005). Insecticidal, repellent and oviposition-deterrent activity of selected essential oils against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. *Bioresource Technology* 96: 1749-1757.
- Rajakumar, S. and Jebanesan, A. (2007). Repellent activity of selected plant essential oils against the malarial fever mosquito *Anopheles stephensi*. *Tropical. Biomedicine* 24:71-75.

- Schreck, C.E. and McGovern, T.P. (1989). Repellants and other personal protection strategies against *Aedes albopictus*. *Journal of the American Mosquito Control Association* 5:247-250.
- Senthil Nathan, S., Chung, P.G. and Murugan, K. (2004). Effect of botanicals and bacterial toxin on the gut enzyme of *Cnaphalocrocis medinalis*. *Phytoparasitica* 32: 433-443.
- Sfara, V., Zerba, E.N. and Alzogaray, R.A. (2009). Fumigant insecticidal activity and repellent effect of five essential oils and seven monoterpenes on first-instar nymphs of *Rhodnius prolixus*. *J Journal of Medical Entomology* 46:511–515.
- Spero, C.N., Gonzalez, I.Y., Scialdone, A.M. and Hallahan, L.D. (2008). Repellency of hydrogenated catmint oil formulations to black flies and mosquitoes in the field. *Journal of Medical Entomology* 45:1080–1086.
- Sritabutra, D., Soonwera, M., Waltanachanobon, S. and Pongjai, S. (2011). Evaluation of herbal essential oil as repellants against *Aedes aegypti* (L.) and *Anopheles dirus* Peyton & Harrion. *Asian Pacific Journal of Tropical Biomedicine*, pp. 5124-5128.
- Tariq, R.M. and Qadri, S.S. (2001). Repellent activity of some local plant's oil, two commercial repellants, di-methyl phthalate and non-alcoholic bitter against dengue vector mosquitoes. *Pakistan Journal of Entomology Karachi* 16: 7-10.
- Traboulsi, A.F., El-Haj, S., Tueni, M., Taoubi, K., Nader, N.B. and Mrad, A. (2005). Repellency and toxicity of aromatic plant extracts against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Management Science* 61: 597-560.
- Trongtokit, Y., Rongsrivam, Y., Komalamisra, N., and Apiwathnasorn, C. (2005). Comparative repellency of essential oils against mosquito bites. *Phytotherapy Research* 19: 303–309.
- Tyagi, B.K., Shahi, A.K. and Koul, B.L. (1998). Evaluation of repellent activities of *Cymbopogon* essential oils against mosquito vectors of malaria, filariasis and dengue fever in India. *Phytomedicine* 5:324-329.
- Venkatachalam, M.R. and Jebanesan, A. (2001). Repellent activity of *Ferronia elephantum* Corr (Rutaceae) leaf extract against *Aedes aegypti* (L). *Bioresource Technology* 76: 287-288.
- WHO, 1995. International travel and health vaccination requirement and health advice. World Health Organization, Geneva.
- WHO, 1996. Report of the WHO informal consultation on the evaluation and testing of insecticides. CTD / WHO PES / IC / 96.1. Geneva: pp. 69.
- WHO, 2009. Guidelines for efficacy testing of mosquito repellants for human skins. WHO/HTM/NTD/WHOPES/2009.WHO:Geneva; p. 4-18.
- Yang, P. and Ma, Y. (2005). Repellent effect of plant essential oil against *Aedes albopictus*. *Journal of Vector Ecology* 30:231–234.

(Received 17 April 2013; accepted 30 June 2013)